

BREAM

SPECIES LIST

BREAM-2

BREAM-2

1000

1500

4/7

2000

2500

3000

3500

4000

4500

5000

5500

4/7

6000

	1000	1500	2000	2500	3000	3500	4000	4500	5000	5500	6000
CALC. BENTHONICS - II											
65. Elphidium imperatrix											
66. E. chapmani	.	.	.	.	.	.	.	.	.	.	.
67. Notorotalia miocenica	.	.	.	.	.	.	.	.	.	.	.
68. Elphidium cf. crassatum	.	.	.	.	.	.	.	.	.	.	.
69. Notorotalia howchini	.	.	.	.	.	.	.	.	.	.	.
70. Elphidium crassatum	.	.	.	.	.	.	.	.	.	.	.
71. E. adalaidense	.	.	.	.	.	.	.	.	.	.	.
CALC. BENTHONICS - III											
72. Carpentaria rotaliformis	.	.	.	.	.	.	.	.	.	.	.
73. Amphistegina lessoni	.	.	.	.	.	.	.	.	.	.	.
CALC. BENTHONIC - IV											
74. Cassidulina laevigata											
75. Nonionella magnalingua	.	.	.	.	.	.	.	.	.	.	.
76. Sphaeroidina bulloides	.	.	.	.	.	.	.	.	.	.	.
77. Chilostomella ovoidea	.	.	.	.	.	.	.	.	.	.	.
78. Pullenia quadraloba	.	.	.	.	.	.	.	.	.	.	.
79. Cassidulinoides orientalis	.	.	.	.	.	.	.	.	.	.	.
80. Cassidulina 'globosa'	.	.	.	.	.	.	.	.	.	.	.
81. C. subglobosa	.	.	.	.	.	.	.	.	.	.	.
82. Pullenia bulloides	.	.	.	.	.	.	.	.	.	.	.
83. Sphaeroidina serrata	.	.	.	.	.	.	.	.	.	.	.
84. Neisocalia subita	.	.	.	.	.	.	.	.	.	.	.
85. Pullenia quinqueloba	.	.	.	.	.	.	.	.	.	.	.
depth	1200	2150			3100		4000	4750	5100	5350	5550
ZONE	A	B		C		D - 2	E	F	G	H	I
											J
											615
											72

