

SIDEWALL CORES
Depth in metres

2505.0	
2514.5	
2525.0	
2534.0	
2537.5	
2540.0	
2541.0	
2542.5	
2543.5	
2548.0	
2549.0	
2550.0	
2551.0	
2552.0	
2555.5	
2561.0	
2562.0	
2567.0	
2568.5	
2573.0	
2575.0	
2583.0	
2700.0	
2703.0	
2716.5	
2726.0	
2733.0	
2744.0	
2751.5	
2759.0	
2786.0	
2787.0	
2854.0	
2881.0	
2924.0	
2943.0	
2982.0	
2991.5	
3002.0	
3003.0	
3017.0	
3027.0	
3036.0	
3047.0	
3055.0	
3231.0	
3332.0	
3368.5	
3382.0 (#8)	
3382.0 (#51)	
3397.0	
3436.0	
3490.0	
3539.0	
3568.0	
3587.0	
3602.0	
3637.5	
3683.5	
3729.0	
3749.0	
3800.0	
3851.5	
3895.0	
3927.5	
3953.0	
3995.5	
4049.0	
4065.0	
4125.0	
4154.0	
4197.0	
4255.5	
4287.5	
4305.5	
4320.0	
4343.0	
4355.0	
4375.0	
4454.0	
4492.0	
4558.0	

Barren Samples (m) Samples not processed (m)
(mudcake)

Reference to Species

1. Dettmann, 1963
2. Dettmann & Playford, 1968
3. Stover & Partridge, 1973
4. Couper, 1960
5. Lentz & Williams, 1977
6. Stover & Evitt, 1978
7. Harris, 1965
8. Foster, 1979
9. Martin, 1973
10. Cookson & Pike, 1953
11. McIntyre, 1965

*Reworked Early Cretaceous. See text for further explanation.
?Identification probable but specimens very poorly preserved or broken up.

Key to assessment of preservation and abundance

- x exceptionally good
- ◊ good, above average
- o average
- ‡ poor, below average
- z exceedingly poor, trace occurrence.

TABLE 1: SPORES, POLLEN and DINOFLAGELLATES IDENTIFIED IN HERMES #1

Helene A Martin, July 1983

W.C.R. W903

DEPT. NAT. RES & ENV

PE900483

