

BASIN GIPPSLAND

DATE _____



WELL NAME KINGFISH -3

ELEVATION +31 feet

AGE	PALYNOLOGIC ZONES	HIGHEST DATA					LOWEST DATA				
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
Eocene	<u>P. tuberculatus</u>										
	<u>U. N. asperus</u>										
	<u>M. N. asperus</u>										
	<u>L. N. asperus</u>										
	<u>P. asperopolus</u>										
	<u>U. M. diversus</u>	7460	1				7482	1			
	<u>M. M. diversus</u>										
	<u>L. M. diversus</u>	7611	2	7620	1		7680	3	7635	2	
Paleocene	<u>U. L. balmei</u>	7900	3	7934	1		8260	1			
	<u>L. L. balmei</u>										
	<u>T. longus</u>										
Cretaceous	<u>T. lilliei</u>										
	<u>N. senectus</u>										
	<u>C. trip./T.pach.</u>										
	<u>C. distocarin.</u>										
	<u>T. pannosus</u>										
EARLY CRETACEOUS											
PRE-CRETACEOUS											
	<u>T.D.</u>	8293									

COMMENTS:

- RATINGS: 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.
 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.
 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.
 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spore and pollen or microplankton, or both.
 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATA RECORDED BY: LES./ADP

DATE June 1971; Dec. 1971

DATA REVISED BY: ADP

DATE Jan. 1975.