

BASIN GIPPSLAND

DATE \_\_\_\_\_

WELL NAME SNAPPER-2

ELEVATION +31 feet



AGE	PALYNOLOGIC ZONES	HIGHEST DATA					LOWEST DATA				
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
Eocene	<u>P. tuberculatus</u>										
	<u>U. N. asperus</u>										
	<u>M. N. asperus</u>										
	<u>L. N. asperus</u>	4036	1				4232	1			
	<u>P. asperopolus</u>	4321	1				4415	1			
	<u>U. M. diversus</u>										
	<u>M. M. diversus</u>										
	<u>L. M. diversus</u>	4862	1				5320	1			
Paleocene	<u>U. L. balmei</u>	5490	1				6608	1			
	<u>L. L. balmei</u>	6732	2				6924	1			
	<u>T. longus</u>	7702	1				8218	1			
Cretaceous	<u>T. lilliei</u>	8620	1	8495	2		8910	1			
	<u>N. senectus</u>										
	<u>C. trip./T.pach.</u>										
	<u>C. distocarini</u>										
	<u>T. pannosus</u>										
EARLY CRETACEOUS											
PRE-CRETACEOUS											
		T.D	10010								

COMMENTS:

Deflandrea heterophlycta Dinoflagellate Zone 4232(2)  
Marine incursions of the Wetz. homomorpha Dinoflagellate Zone  
occurs only in the two samples at 6144' and 6608'

RATINGS:

- 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.
- 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.
- 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.
- 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spore and pollen or microplankton, or both.
- 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATA RECORDED BY: LES./ADP

DATE June 1971; Dec. 1971.

DATA REVISED BY: ADP

DATE Jan. 1975.