

BASIN GIPPSLAND

DATE \_\_\_\_\_

WELL NAME TAILOR-1

ELEVATION + 31 FEET



AGE	PALYNOLOGIC ZONES	HIGHEST DATA				LOWEST DATA					
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
Eocene	<u>P. tuberculatus</u>										
	<u>U. N. asperus</u>										
	<u>M. N. asperus</u>										
	<u>L. N. asperus</u>	7896	2				7896	2			
	<u>P. asperopolus</u>										
	<u>U. M. diversus</u>										
	<u>M. M. diversus</u>										
	<u>L. M. diversus</u>	7918	1				7918	1			
Paleocene	<u>U. L. balmei</u>	7941	0				8318	0			
	<u>L. L. balmei</u>										
	<u>T. longus</u>										
Cretaceous	<u>T. lilliei</u>										
	<u>N. senectus</u>										
	<u>C. trip./T.pach.</u>										
	<u>C. distocarin.</u>										
	<u>T. pannosus</u>										
EARLY CRETACEOUS											
PRE-CRETACEOUS											
T.D.		8498									

COMMENTS: Wetzeliella homomorpha Dinoflagellate Zone 7941 (0) - 8318(0)

- RATINGS: 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.  
 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.  
 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.  
 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spore and pollen or microplankton, or both.  
 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATA RECORDED BY: LES/ADP

DATE June 1971; Dec. 1971.

DATA REVISED BY: ADP

DATE Jan. 1975.