



WASH. STATE UNIVERSITY
 LOCAL NAME: SALMON ELEVATION: 199 feet
 DATE: June 1971

AGE	PALYNOLOGIC ZONES	HIGHEST DATA				LOWEST DATA					
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
C. REC.	<i>T. helios</i>										
	<i>P. tuberculatus</i>										
Eocene	<i>U. R. asperus</i>										
	<i>L. R. asperus</i>	6595	2			6710	4				
	<i>P. asperonolus</i>	6888	1			6888	1				
	<i>U. H. diversus</i>	7040	1			7310	2	7220	1		
	<i>L. H. diversus</i>	7640	2			7544	2				
FALES-GENE	<i>L. ballasi</i>	8008	1			9250	2				
	<i>T. longus</i>										
LATE CRETACEOUS	<i>T. hillii</i>										
	<i>H. senectus</i>										
	<i>C. trip./T. pach.</i>										
	<i>C. distocarin.</i>										
	<i>T. pannosus</i>										
EARLY CRETACEOUS	<i>C. paradoxa</i>										
	<i>C. striatus</i>										
	<i>U. C. hughesii</i>										
	<i>L. C. hughesii</i>										
	<i>C. stylosus</i>										
Pre-Cretaceous											

COMMENTS: Samples below 8008 feet contain too few specimens for reliable interpretation; additional processing required.

T.D. 9855 (1966)

- RATINGS: 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.
 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.
 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.
 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spores and pollen or microplankton, or both.
 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATE RECORDED BY: L.E. Stover / A.D. Partridge DATE: June 1971
 DATA REVIEWED BY: CHECKED: L.E.S. DATE: Dec. 1971