

APPENDIX-5

PALYNOLOGICAL ANALYSIS, TARWHINE-1,
GIPPSLAND BASIN

by

M.K. Macphail

INTRODUCTION:

Seventy six (76) sidewall cores and two core samples were processed and examined for spore-pollen. The recovery of microfossils was good to fair for most samples except in the Lygistepollenites balmei Zone. Preservation was usually poor and relatively few samples contained zone species of spores and pollen. Age determinations are supported by an analysis of the dinoflagellate assemblages by A.D. Partridge.

Palynological zones and lithological facies subdivisions from the base of the Lakes Entrance Formation to the total depth of the well are given below. Table 1 represents a summary of the palynological analyses. The occurrence of each spore and pollen type is tabulated in the accompanying range chart.

SUMMARY

<u>UNIT/FACIES</u>	<u>ZONE</u>	<u>DEPTH (metres)</u>
LAKES ENTRANCE FORMATION	<u>P. tuberculatus</u>	1310
GURNARD FORMATION	Upper <u>N. asperus</u>	1345-1348
	Middle <u>N. asperus</u>	1351-1381
	Middle <u>N. asperus</u>	1392-1405
	Lower <u>N. asperus</u>	1412-1591
	<u>P. asperopolus</u>	1643-1656
LATROBE GROUP COARSE CLASTICS	Upper <u>M. diversus</u>	1666-1727
	Middle to Lower	1818-1892
	<u>M. diversus</u>	
	Upper <u>L. balmei</u>	1911-1932
	Lower <u>L. balmei</u>	2026-2391
	<u>T. longus</u>	2420-2934
	<u>T. lilliei</u>	2940-2949

GEOLOGICAL COMMENTS:

1. The Tarwhine-1 well contains a continuous sequence of zones from the Late Cretaceous T. lilliei Zone to the Early Oligocene P. tuberculatus Zone. Although spore-pollen assemblages were not sufficiently diverse to separate the Middle and Lower M. diversus Zones, it is probable both are represented in the well.

2. The Gurnard Formation, recognised by glauconitic sediments from 1345 to 1386 metres, contains both Middle and Upper N. asperus spore-pollen and dinoflagellate assemblages. The Upper N. asperus Zone is represented in sidewall cores at 1345 and 1348 metres. The lower of the two also contains foraminiferal faunas referable to the Late Eocene Zone K (Hannah 1982). The Gurnard Formation, initially picked on cutting lithology at the time of drilling as occurring between 1370 to 1386 metres, is clearly Middle N. asperus in age.

3. The Middle N. asperus Zone extends down into the coarse clastics below the Gurnard Formation, demonstrating that there is no significant unconformity at this lithological boundary. Dinoflagellates remain common across the boundary to 1412 metres, top of the Lower N. asperus Zone. This is close to the uppermost occurrence of coal at 1407 metres and consistent with a change from a terrestrial/fluviial to an open marine environment during the Middle Eocene.

4. The presence of Upper N. asperus age sediments in the Tarwhine-1 well strengthens the probability (Partridge 1977) that the unsampled interval between 1040 to 1045 metres in

the Barracouta-4 well is also Upper N. asperus in age, i.e., that there is no unconformity between the Middle N. asperus Gurnard Formation and the P. tuberculatus Lakes Entrance Formation at Barracouta-4. A similar sequence of time-rock units at the two well sites is consistent with a Middle Eocene shoreline parallel to the present-day coastline.

5. Although no major periods of non-deposition and/or erosion are evident, mean sedimentation rates show a sustained decrease from greater than 80 metres per million years in the T. longus Zone to less than 30 metres per million years in the Middle N. asperus Zone. This may reflect a shift (northwards) in the depositional centre of the basin away from the Tarwhine area. Anomalously low deposition rates calculated for the Upper N. asperus and P. asperopolus Zones suggest that minor periods of erosion/non-deposition have occurred. Alternatively, portions of the undated interval between 1644 and 1591 metres may also be P. asperopolus in age.
6. There is no evidence in the form of dinoflagellate cysts for any marine transgression at the Tarwhine-1 well site before the top of the Lower N. asperus Zone at 1412 metres.
7. The Thermal Alteration Index for all samples is less than 2.

DISCUSSION OF ZONES.

Zone boundaries have been established using the criteria of Stover and Evans (1973), Stover and Partridge (1973) and subsequent revisions (A.D. Partridge per comm.).

Proteacidites tuberculatus Zone - above 1310 metres.

The zone is recognised on the presence of Cyatheacidites annulatus at 1250 metres and 1310 metres. The other sample at 1280 metres contains Quintinia psilatispora Martin. This is the lowest stratigraphical record of this taxon in the Gippsland Basin to date.

Upper Nothofagidites asperus Zone: 1345 - 1348 metres

This zone is identified by the absence of spore-pollen indicator species of the underlying zone (e.g., Triorites magnificus) and the overlying zone (e.g., Cyatheacidites annulatus). The age-determination is supported by (i) Pthanoperidinium comatum being the dominant dinoflagellate species in both samples, (ii) Proteacidites rectomarginis being the most common large proteaceous pollen and (iii) Nothofagidites flemingii being a common element in the lowermost sample.

Middle Nothofagidites asperus Zone and Vozzkenikova extensa

Dinoflagellate Zone: 1351.1 - 1405 metres.

The top of the zone is based on the highest occurrence of the pollen taxa Triorites magnificus and Proteacidites adenanthoides, and the dinoflagellates V. extensa and Corrudinium incompositum, at 1351.1 metres. The base of the zone is placed at the lowest occurrence of V. extensa, in samples at 1405.44 metres (core) and 1401.8 metres (sidewall core). Spore-pollen diagnostic of the zone appear later in the zone: Anacolosidites sectus at 1392 metres and Triorites magnificus at 1380.9 metres (base of the Gurnard Formation). The late appearance of diagnostic species is attributed to poor concentration and poor preservation of spore-pollen in samples from the hydrocarbon column at the top of the coarse clastics. Although these samples are rich in terrestrially-derived organic matter, oxidation tends to destroy all forms of organic material except for the highly resistant dinoflagellate cysts.

Lower Nothofagidites asperus Zone: 1411.8 - 1591 metres.

The base of the zone is placed at the marked increase in Nothofagidites spp. abundance, from less than 1% at 1643.5 metres to 33% at 1591 metres. Nothofagidites asperus is recorded at the base but other zone species are recorded higher within the zone, e.g., N. falcatus at 1426.6 metres, and Rugulatisporites trophus at 1542.4 metres. The top of the zone is picked by the absence of indicator species of the overlying zone, i.e., Triorites magnificus and V. extensa. The two lowermost samples, at 1542.4 metres and 1591 metres, contain abundant Proteacidites asperopolus pollen. This phenomenon has been previously noted for Lower N. asperus age samples in the Barracouta-4 well (Partridge 1977).

Proteacidites asperopolus Zone: 1643.5 - 1656 metres.

Two sidewall core samples are identified as belonging to this zone. Confidence in the age-determination of the uppermost, at 1643.5 metres, is good due to the first appearance here of Sapotaceoidaepollenites rotundus, the occurrence of Myrtacidites tenuis, a species which ranges no higher than the P. asperopolus Zone, and the dominance of proteaceous pollen, particularly P. pachyopolus, P. reticuloscabratus and P. xestiformis. Although none of these species is restricted to the P. asperopolus Zone, the marked difference in composition between these samples, the overlying Nothofagidites spp. -dominant and underlying Malvacipollis -dominant palynofloras is typical of the zone. The lower of the two samples, 1656 metres, contains M. tenuis. The remainder of this spore-pollen assemblage consists of long-ranging taxa and the sample is assigned a P. asperopolus age on the basis that it is more similar to sample 1643.5 metres than those in the underlying M. diversus Zone.

Upper Malvacipollis diversus Zone: 1666.4 - 1727 metres.

The top of the zone is identified by the first occurrence of abundant M. diversus pollen and also contains the first appearance of Myrtaceidites tenuis. The occurrence of abundant M. diversus, Proteacidites pachypolus and Milfordia homeopunctata pollen confirm an Upper M. diversus age for samples at 1715.5 and 1727 metres. The quartz-rich sediments between 1727 and 1818.5 metres were barren or contained reworked spore-pollen only. e.g., Gambierina rudata.

Middle-Lower Malvacipollis diversus Zone: 1818.5 - 1892.5 metres.

The base of the zone is picked at the first occurrence of M. diversus pollen, at 1892.5 metres. The species is relatively common in this sample and at 1882.1 metres but is absent from samples between 1818.5 and 1882.1 metres. The top of the zone, at 1818.5 metres is based on the absence of indicator species of the Upper M. diversus Zone, e.g., Proteacidites pachypolus. However this sample cannot be definitely assigned to the Middle M. diversus Zone due to the absence of Middle M. diversus indicator species such as Proteacidites ornatus, P. tuberculiformis and P. xestiformis. The variable occurrences of Cupanieidites orthoteichus, Ischyosporites irregularis, Proteacidites lapis, P. obscurus, Tricolporites adelaidensis, T. paenestriatus and Triporopollenites helosus demonstrate the interval is no older than M. diversus age but the low yields and generally poor spore-pollen diversity make it impossible to make a more detailed subdivision.

Upper Lygistepollenites balmei Zone: 1911 - 1932 metres.

The top of the zone is identified by the first appearance of L. balmei (abundant) with Verrucosisporites kopukuensis. The presence of Gambierina edwardsii, Nothofagidites endurus and Phyllocladidites verrucosus demonstrate the sample is no younger

The top of the zone is defined by the occurrence of Grapnelispora evansii sp. nov. at 2420 metres. The highest occurrence of T. longus is at 2431 metres. Quadruplanus brossus which is diagnostic of the zone occurs lower down, at 2445 metres. The base of the zone is picked on the simultaneous first appearances of T. longus, Proteacidites otwayensis and P. reticuloconcavus at 2934.5 metres. There is currently some uncertainty as to the exact stratigraphic range of these Palaeocene Proteacidites spp. and subsequent work may show the first appearance of Stereisporites (Tripunctisporis) punctatus at 2699.3 metres represents the base of the T. longus Zone.

Tricolporites lillieii Zone: 2939.8 - 2948.5 metres.

The zone is characterised by a marked increase in Nothofagidites endurus relative to other spore-pollen species (Stover & Evans 1973). These authors distinguish the base of the overlying T. longus Zone by the virtual absence of Nothofagidites spp. and a marked increase in Triorites edwardsii (= Gambierina rudata) pollen. On these criteria, the age of the sediment at 2934.5 metres is equivocal since G. rudata is abundant and N. endurus frequent.

The top of the zone is picked at 2939.8 metres, the highest sample in which N. endurus is clearly more abundant than G. rudata. T. lillieii is also present. The basal sample, at 2948.5 metres, contains a mixed late Cretaceous, Paleocene and Eocene spore-pollen assemblage, lacking T. lillieii. The sample has been contaminated by cavings but is obviously no older than the T. lillieii Zone due to the presence of G. rudata and Tripoporopollenites sectilis.

than Upper L. balmei in age. V. kopukuensis at 1926 metres confirms an Upper L. balmei age for this sample. The base of the zone is picked at 1932 metres due to, (i) the absence of indicator species of the Lower L. balmei Zone, e.g., Tetracolporites verrucosus, and (ii) an abundance of Australopollis obscurus. Occurrences of Drytopollenites semilunatus at 1911 metres and 1932 metres and Periporopollenites demarcatus at 1932 metres and 2496.1 metres represent extensions in the range of these taxa into the Paleocene and late Cretaceous respectively.

Lower Lygistepollenites balmei Zone: 2026.6 - 2391.5 metres.

The zone falls within an interval in which most samples are either barren or contain spore-pollen assemblages of low diversity and limited stratigraphic usefulness. The top of the zone, 2026.6 metres, is identified by the highest occurrence of Tetracolporites verrucosus and the first occurrence of Proteacidites annularis. The base of the zone is picked at 2391.5 metres on the lowermost occurrence of an L. balmei and (abundant) Gambierina rudata assemblage lacking indicator species of the T. longus Zone. Spore-pollen diagnostic of sediments no older than L. balmei were recorded higher in the zone, e.g., Basopollis mutabilis at 2346 metres, Haloragacidites harrisii and Malvacipollis subtilis at 2036 metres.

Tricolpites longus Zone: 2420 - 2934.5 metres.

The zone is characterised by spore-pollen assemblages that are more diverse than in the overlying L. balmei Zone. Gambierina rudata and gymnosperms are the dominant taxa but many samples also contain T. longus and species which first appear in the T. longus zone, e.g., Tetracolporites verrucosus. Triporopollenites sectilis and Tricolporites lilliei occur in most samples.

R E F E R E N C E S

- HANNAH, M.J., 1982. Micropalaeontological analysis of Tarwhine-1, Gippsland Basin, Victoria. Esso Australia Ltd, Palaeontological Report: 1982/18.
- PARTRIDGE, A.D., 1977. Palynological Analysis Barracouta-4, Gippsland Basin. Esso Australia Ltd, Palaeontological Report: 1977/16.
- STOVER, L.E., & EVANS, P.R., 1973. Upper Cretaceous spore-pollen zonation, offshore Gippsland Basin, Australia. Special Publication Geological Society of Australia 4: 55-72.
- STOVER, L.E., & PARTRIDGE, A.D., 1973. Tertiary and Late Cretaceous spores and pollen from the Gippsland Basin, Southeastern Australia. Proceedings of the Royal Society of Victoria, 85: 237-86.

BASIC DATA

Table-1: Palynological Data
Range Chart - Dinoflagellates
Range Chart - Spore Pollen

P A L Y N O L O G Y D A T A S H E E T

B A S I N : GIPPSLAND
WELL NAME : TARWHINE-1

ELEVATION: KB: 21m GL: -48m
TOTAL DEPTH: 2955 metres

A G E	PALYNOLOGICAL ZONES	H I G H E S T D A T A					L O W E S T D A T A				
		Preferred Depth	Rtg	Alternate Depth	Rtg	Two Way Time	Preferred Depth	Rtg	Alternate Depth	Rtg	Two Way Time
NEOGENE	<i>T. pleistocenicus</i>										
	<i>M. lipsis</i>										
	<i>C. bifurcatus</i>										
	<i>T. bellus</i>										
PALEOGENE	<i>P. tuberculatus</i>	1250.0	0				1310.0	0			
	Upper <i>N. asperus</i>	1345.0	1				1348.0	1			
	Mid <i>N. asperus</i>	1351.1	0				1405.4	2	1392.0	1	
	Lower <i>N. asperus</i>	1411.8	2				1591.0	2	1542.4	1	
	<i>P. asperopolus</i>	1643.5	2				1656.0	2			
	Upper <i>M. diversus</i>	1666.4	1				1727.0	1			
	Mid <i>M. diversus</i>	1818.5	2								
	Lower <i>M. diversus</i>						1892.5	1			
	Upper <i>L. balmei</i>	1911.0	0				1932.0	2	1926.0	0	
	Lower <i>L. balmei</i>	2026.6	1				2391.5	2	2346.0	1	
LATE CRETACEOUS	<i>T. longus</i>	2420.0	0				2934.5	2	2699.3	1	
	<i>T. lilliei</i>	2939.8	1				2948.5	2	2939.8	1	
	<i>N. senectus</i>										
	U. <i>T. pachyexinus</i>										
	L. <i>T. pachyexinus</i>										
	<i>C. triplex</i>										
EARLY CRET.	<i>A. distocarinatus</i>										
	<i>C. paradoxus</i>										
	<i>C. striatus</i>										
	<i>F. asymmetricus</i>										
	<i>F. wonthaggiensis</i>										
	<i>C. australiensis</i>										
	PRE-CRETACEOUS										

COMMENTS: It has not been possible to subdivide the Middle and Lower *M. diversus* Zones due to poor diversity and poor preservation of spore-pollen.

- CONFIDENCE RATING:
- 0: SWC or Core, Excellent Confidence, assemblage with zone species of spores, pollen and microplankton.
 - 1: SWC or Core, Good Confidence, assemblage with zone species of spores and pollen or microplankton.
 - 2: SWC or Core, Poor Confidence, assemblage with non-diagnostic spores, pollen and/or microplankton.
 - 3: Cuttings, Fair Confidence, assemblage with zone species of either spores and pollen or microplankton, or both.
 - 4: Cuttings, No Confidence, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If an entry is given a 3 or 4 confidence rating, an alternative depth with a better confidence rating should be entered, if possible. If a sample cannot be assigned to one particular zone, then no entry should be made, unless a range of zones is given where the highest possible limit will appear in one zone and the lowest possible limit in another.

DATA RECORDED BY: M.K. Macphail & A.D. Partridge DATE: May 4, 1982.

DATA REVISED BY: _____ DATE: _____

TABLE-1
PALYNOLOGICAL DATA
BASIC DATA

SAMPLE NO.	DEPTH (Metres)	YIELD	DIVERSITY SPORE POLLEN
SWC 139	1250	Good	Low
SWC 138	1280	Good	Low
SWC 137	1310	Good	Low
SWC 135	1345	Good	Moderate
SWC 134	1348	Good	High
SWC 133	1351.1	Good	High
SWC 132	1353	Good	High
SWC 130	1359	Good	Moderate
SWC 127	1367.1	Fair	Moderate
SWC 123	1375.1	Good	High
SWC 120	1380.9	Good	Moderate
SWC 116	1392	Good	Moderate
SWC 113	1401	Good	High
Core 2	1405.44	Good	Low
Core 2	1411.8	Good	Moderate
SWC 101	1422.1	Low	Poor
SWC 99	1426.6	Good	Moderate
SWC 92	1503	Good	Moderate
SWC 89	1542.4	Good	Moderate
SWC 86	1591	Good	Moderate
SWC 82	1643.5	Good	Moderate
SWC 81	1656	Low	Low
SWC 107	1666.4	Good	Moderate
SWC 78	1693	Nil	-
SWC 76	1715.5	Fair	Moderate
SWC 75	1727	Good	High
SWC 73	1754	Nil	-
SWC 69	1807	Nil	-
SWC 68	1818.5	Good	Moderate
SWC 67	1836	Good	Moderate
SWC 65	1860	Fair	Low
SWC 64	1882.1	Low	Low
SWC 62	1892.5	Low	Low
SWC 61	1900	Very Low	Very Low
SWC 60	1911	Good	High
SWC 59	1926	Fair	Low
SWC 58	1932	Good	Moderate
SWC 57	1955	Nil	-
SWC 51	2003.1	Very Low	Very Low
SWC 49	2026.6	Good	Moderate
SWC 48	2036	Fair	Moderate
SWC 41	2115.7	Fair	Low
SWC 38	2163	Fair	Low
SWC 36	2196.2	Nil	-
SWC 35	2204.9	Fair	Low
SWC 33	2230.9	Nil	-
SWC 32	2241.1	Very Low	Very Low
SWC 31	2255.1	Nil	-
SWC 29	2273	Nil	-
SWC 28	2290.1	Very Low	Very Low
SWC 27	2304	Low	Low
SWC 26	2315	Fair	Low
SWC 24	2346	Fair	Low
SWC 23	2352.5	Nil	-

TABLE-1
SUMMARY OF PALYNOLOGICAL ANALYSIS, TARWHINE-1, GIPPSLAND BASIN
INTERPRETATIVE DATA

SAMPLE NO.	DEPTH (Metres)	YIELD	DIVERSITY SPORE POLLEN	LITHOLOGY	ZONE	AGE	CONFIDENCE RATING	COMMENTS
SWC 139	1250	Good	Low	Marl	<u>P. tuberculatus</u>	Oligocene	0	<u>Cyatheacidites annulatus</u>
SWC 138	1280	Good	Low	Marl	<u>P. tuberculatus</u>	Oligocene	0	<u>Quintinia psilatospora</u>
SWC 137	1310	Good	Low	C. lut.	<u>P. tuberculatus</u>	Oligocene	0	<u>Cyatheacidites annulatus</u>
SWC 135	1345	Good	Moderate	Clyst, calc, glauc.	<u>Upper N. asperus</u>	Late Eocene	2	Diverse <u>Nothofagidites</u> assemblage
SWC 134	1348	Good	High	Clyst, calc, glauc.	<u>Upper N. asperus</u>	Late Eocene	1	<u>Proteacidites rectomarginis</u> and <u>Phthanoperidinium</u> common.
SWC 133	1351.1	Good	High	Clyst, calc.	<u>Middle N. asperus</u>	Late Eocene	0	<u>Triorites magnificus</u>
SWC 132	1353	Good	High	Clyst, calc.	<u>Middle N. asperus</u>	Late Eocene	0	<u>Triorites magnificus</u> , <u>Aglaotheidis qualumis</u> , <u>Proteacidites pachypolus</u> , <u>Simplicipollis meridianus</u>
SWC 130	1359	Good	Moderate	Clyst, tr. glauc.	<u>Middle N. asperus</u>	Late Eocene	2	<u>Proteacidites rectomarginis</u> common
SWC 127	1367.1	Fair	Moderate	Clyst, calc. tr. glauc.	<u>Middle N. asperus</u>	Late Eocene	2	<u>Rugulatisporites trophus</u>
SWC 123	1375.1	Good	High	Clyst, calc, glauc.	<u>Middle N. asperus</u>	Late Eocene	0	<u>Triorites magnificus</u>
SWC 120	1380.9	Good	Moderate	Stst, glauc.	<u>Middle N. asperus</u>	Late Eocene	1	<u>Aglaotheidia qualumis</u> , <u>Tripoporollenites ambiguus</u> , <u>Proteacidites recavus</u>
SWC 116	1392	Good	Moderate	Sltst, calc, carb.	<u>Middle N. asperus</u>	Late Eocene	1	<u>Anacolosidites sectus</u> , <u>Proteacidites crassus</u>
SWC 113	1401	Good	High	Sst, clay-rich	<u>Middle N. asperus</u>	Late Eocene	1	<u>Dryptopollenites semilunatus</u> , <u>Schizocolpus marlinensis</u> , <u>Proteacidites crassus</u>
Core 2	1405.44	Good	Low	---	<u>Middle N. asperus</u>	Middle Eocene	2	<u>Malvacipollis robustus</u> . Sample from oil column.
Core 2	1411.8	Good	Moderate	---	<u>Lower N. asperus</u>	Middle Eocene	2	<u>Malvacipollis robustus</u> , <u>Proteacidites pachypolus</u> , Sample from oil column.
SWC 101	1422.1	Low	Poor	---	---	---	-	---
SWC 99	1426.6	Good	Moderate	Sst, carb. lens	<u>Lower N. asperus</u>	Middle Eocene	2	<u>Nothofagidites falcatus</u>
SWC 92	1503	Good	Moderate	Sst, carb.	<u>Lower N. asperus</u>	Middle Eocene	2	<u>Proteacidites asperopolus</u> , <u>P. pachypolus</u>

TABLE-1 Cont. 2

SAMPLE NO.	DEPTH (Metres)	YIELD	DIVERSITY		LITHOLOGY	ZONE	AGE	CONFIDENCE RATING	COMMENTS
			SPORE	POLLEN					
SWC 89	1542.4	Good	Moderate	Clyst		Lower <u>N. asperus</u>	Middle Eocene	1	<u>Rugulatisporites trophus</u>
SWC 86	1591	Good	Moderate	Clyst		Lower <u>N. asperus</u>	Middle Eocene	0	<u>Rugulatisporites trophus</u> , <u>Proteacidites asperopolus</u> & <u>P. rugulatus</u> common
SWC 82	1643.5	Good	Moderate	Clyst, carb.		<u>P. asperopolus</u>	Middle Eocene	1	<u>Sapotaceoidaepollenites rotundus</u> , <u>Myrtacidites tenuis</u> , abundant <u>Proteacidites</u>
SWC 81	1656	Low	Low	Sst, carb. lens		Indeterminate	---	-	<u>Nothofagidites</u> spp. rare, <u>Myrtacidites tenuis</u>
SWC 107	1666.4	Good	Moderate	Sst, carb. stringers		Upper <u>M. diversus</u>	Early Eocene	1	<u>M. diversus</u> dominant; <u>Proteacidites pachypolus</u>
SWC 78	1693	Nil	-	Clyst		Indeterminate	---	-	---
SWC 76	1715.5	Fair	Moderate	Ss		Upper <u>M. diversus</u>	Early Eocene	2	<u>M. diversus</u> dominant, <u>P. pachypolus</u>
SWC 75	1727	Good	High	Slst		Upper <u>M. diversus</u>	Early Eocene	1	<u>Proteacidites pachypolus</u> , <u>P. tuberculiformis</u>
SWC 73	1754	Nil	-	Sst		Indeterminate	---	-	---
SWC 69	1807	Nil	-	Clyst		Indeterminate	---	-	---
SWC 68	1818.5	Good	Moderate	Sst, carb. lens		Middle-Lower <u>M. diversus</u>	Early Eocene	1	<u>Tricolporites moultonii</u>
SWC 67	1836	Good	Moderate	Clyst		M-L <u>M. diversus</u>	Early Eocene	2	<u>Proteacidites lapis</u>
SWC 65	1860	Fair	Low	Sltst		M-L <u>M. diversus</u>	Early Eocene	2	<u>Proteacidites leightonii</u>
SWC 64	1892.1	Low	Low	Sst, carb.		M-L <u>M. diversus</u>	Early Eocene	2	Infrequent <u>M. diversus</u>
SWC 62	1892.5	Low	Low	Sst		M-L <u>M. diversus</u>	Early Eocene	1	<u>M. diversus</u> frequent.
SWC 61	1900	Very Low	Very Low	Sst, carb. lens		Indeterminate	---	-	---
SWC 60	1911	Good	High	Sst, carb.		Upper <u>L. balmei</u>	Late Paleocene	0	<u>L. balmei</u> frequent
SWC 59	1926	Fair	Low	---		Upper <u>L. balmei</u>	Late Paleocene	0	<u>L. balmei</u> frequent <u>Verrucosiporites kopukuensis</u>
SWC 58	1932	Good	Moderate	Clyst		Upper <u>L. balmei</u>	Late Paleocene	2	<u>Australopollis obscurus</u> abundant, <u>L. balmei</u> , <u>Polycolpites langstonii</u>
SWC 57	1955	Nil	-	---		Indeterminate	---	-	---
SWC 51	2003.1	Very Low	Very Low	---		Indeterminate	---	-	<u>Haloragacidites harrisii</u>
SWC 49	2026.6	Good	Moderate	Ss - Slst		Lower <u>L. balmei</u>	Early-Middle Paleocene	1	<u>Tetracolporites verrucosus</u> , <u>Proteacidites annularis</u>

TABLE-1 Cont. 3

SAMPLE NO.	DEPTH (Metres)	YIELD	DIVERSITY		LITHOLOGY	ZONE	AGE	CONFIDENCE RATING	COMMENTS
			SPORE	POLLEN					
SWC 48	2036	Fair	Moderate	Sltst	Lower <u>L. balmei</u>	Early-Middle Paleocene	1	<u>Tetracolporites verrucosus</u>	
SWC 41	2115.7	Fair	Low	Clyst	Lower <u>L. balmei</u>	Early-Middle Paleocene	2	<u>L. balmei</u> , <u>Nothofagidites endurus</u> common	
SWC 38	2163	Fair	Low	Shale	Lower <u>L. balmei</u>	Early-Middle Paleocene	2	<u>L. balmei</u> common	
SWC 36	2196.2	Nil	-	Shale	Indeterminate	---	-	---	
SWC 35	2204.9	Fair	Low	Shale	<u>L. balmei</u>	Paleocene	2	<u>Australopollis obscurus</u> abundant	
SWC 33	2230.9	Nil	-	Clyst	Indeterminate	---	-	---	
SWC 32	2241.1	Very Low	Very Low	Ss	Indeterminate	---	-	---	
SWC 31	2255.1	Nil	-	Shale	Indeterminate	---	-	---	
SWC 29	2273	Nil	-	Sltst	Indeterminate	---	-	---	
SWC 28	2290.1	Very Low	Very Low	Sltst	Indeterminate	---	-	---	
SWC 27	2304	Low	Low	Clyst	Indeterminate	---	-	<u>Latrobosporites cf. ohaiensis</u>	
SWC 26	2315	Fair	Low	Clyst	Lower <u>L. balmei</u>	Early-Middle Paleocene	2	<u>Tetracolporites verrucosus</u> frequent	
SWC 24	2346	Fair	Low	Sltst	<u>L. balmei</u>	Paleocene	1	<u>Basopollis mutabilis</u>	
SWC 23	2352.5	Nil	-	Ss	Indeterminate	---	-	Amorphous substrates	
SWC 20	2362	Low	Very Low	Ss, carb. fragments	Indeterminate	---	-	<u>Gambierina edwardsii</u>	
SWC 17	2374	Low	Low	Clyst	<u>L. balmei</u>	Paleocene	2	<u>Gambierina rudata</u> frequent, <u>Stereisporites</u> (<u>Tripunctisporis</u>) <u>punctatus</u>	
SWC 13	2391.5	Low	Low	Clyst	<u>L. balmei</u>	Paleocene	2	<u>Gambierina rudata</u> (common), <u>L. balmei</u>	
SWC 11	2401.2	Nil	-	Clyst	Indeterminate	---	-	Amorphous substrates	
SWC 10	2420	Low	Moderate	Clyst	<u>T. longus</u>	Late Cretaceous	0	<u>Craonelispora evansii</u>	
SWC 9	2431	Good	Moderate	Shale	<u>T. longus</u>	Late Cretaceous	1	<u>T. longus</u> , <u>Tetracolporites verrucosus</u>	
SWC 52	2445	Good	High	--	<u>T. longus</u>	Late Cretaceous	0	<u>Quadrplanus brossus</u>	
SWC 7	2457.2	Fair	Low	Sltst, carb.	<u>T. longus</u>	Late Cretaceous	0	<u>Quadrplanus brossus</u>	
SWC 6	2465.2	Fair	Very Low	Finely bdd argil. mtx	Indeterminate	---	-	<u>Gambierina rudata</u> only.	
SWC 3	2496.1	Fair	Low	Clyst	<u>T. longus</u>	Late Cretaceous	2	<u>Tetracolporites verrucosus</u>	
SWC 245	2571.8	Fair	Low	Ss, trace coal	<u>T. longus</u>	Late Cretaceous	0	<u>Quadrplanus brossus</u> with abundant <u>Gambierina rudata</u>	
SWC 240	2608.3	Fair	Very Low	Sltst	Indeterminate	---	-	<u>Tricolporites lilliei</u>	

TABLE-1 Cont. 4

SAMPLE NO.	DEPTH (Metres)	YIELD	DIVERSITY		LITHOLOGY	ZONE	AGE	CONFIDENCE		COMMENTS
			SPORE	POLLEN				RATING		
SWC 232	2646.5	Fair	Moderate		Slst, some coal	<u>T. longus</u>	Late Cretaceous	1		<u>T. longus</u> , <u>Proteacidites reticuloconcaus</u> , <u>Stereisporites (Tripunctisporis) punctatus</u>
SWC 217	2699.3	Fair	Low		Slst, carb.	<u>T. longus</u>	Late Cretaceous	1		Lowest occurrence of <u>Stereisporites (Tripunctisporis) punctatus</u>
SWC 194	2776	Fair	Low		Ss, some coal	<u>T. longus</u>	Late Cretaceous	2		<u>Proteacidites reticuloconcaus</u> , <u>Tricolporites lilliei</u> , common <u>Gambierina rudata</u>
SWC 187	2799	Fair	Moderate		Ss, some coal	<u>T. longus</u>	Late Cretaceous	2		<u>Proteacidites reticuloconcaus</u> , <u>P. analoxinus</u> , <u>P. angulatus</u>
SWC 185	2807	Very Low	Very Low		Ss	Indeterminate	---	-		---
SWC 177	2858.5	Good	Moderate		Ss	<u>T. longus</u>	Late Cretaceous	2		<u>Proteacidites otwayensis</u> , <u>Tetracolporites verrucosus</u>
SWC 175	2868.4	Very Low	Very Low		Ss	Indeterminate	---	-		Mud contamination
SWC 166	2901.5	Fair	Moderate		Sltst	Indeterminate	---	-		<u>Gambierina rudata</u> common, <u>Tricolporites lilliei</u>
SWC 157	2934.5	Good	Moderate		Slst	<u>T. longus</u>	Late Cretaceous	2		<u>Proteacidites otwayensis</u> , <u>P. reticuloconcaus</u> , <u>T. longus</u> , <u>Gambierina rudata</u> common, <u>Nothofagidites endurus</u> frequent
SWC 156	2939.8	Low	Low		Sltst	<u>T. lilliei</u>	Late Cretaceous	1		<u>Nothofagidites endurus</u> , <u>N. senectus</u> more common than <u>Gambierina rudata</u> , <u>T. lilliei</u>
SWC 154	2948.5	Fair	Moderate		Ss	<u>T. lilliei</u>	Late Cretaceous	2		Contaminated sample but no older than <u>T. lilliei</u> due to occurrence of <u>Gambierina rudata</u> , <u>Triporopollenites sectilis</u>