

11 APR 1991

**PALYNOLOGY AND SOURCE ROCK STUDY
OF PEP 109 AND PEP 110
ONSHORE GIPPSLAND BASIN**

Cluff Resources Pacific Limited
Level 5, 53 Walker Street
North Sydney NSW 2060

April, 1991

**PALYNOLOGY AND SOURCE ROCK STUDY
OF PEP 109 AND PEP 110
ONSHORE GIPPSLAND BASIN**

Cluff Resources Pacific Limited
Level 5, 53 Walker Street
North Sydney NSW 2060

April, 1991

CONTENTS

1. SUMMARY AND CONCLUSIONS

2. APPENDICES:

Appendix 1

Palynology of 23 Subsurface Samples, Onshore Gippsland Basin
by Roger Morgan

Appendix 2

Report on Source Rock Study on 36 Samples from Onshore
Gippsland by Amdel Core Services

3. LIST OF ENCLOSURES

Enclosure 1: Cross Section - Lake Wellington Park to
Merriman-1

Enclosure 2: Cross Section - Golden Beach - 1A to Colliers
Hill

1. SUMMARY AND CONCLUSION

In order to clarify the ambiguity about the presence of Late Cretaceous Strata, equivalent to the Golden beach Formation and confirmation on presence of the Strzelecki Group Sequence in few wells, Cluff Resources Pacific submitted 23 samples for palynological study from 7 wells in the onshore Gippsland Basin, namely Burong-1, Carrs Creek-1, Dutson Downs-1, Merriman-1, Seaspray-1, North Seaspray-1 and Wellington Park-1.

Also 36 core and cutting samples from 6 wells were submitted to Amdel Core Services for geochemical analysis to determine the source potential and maturity f the Latrobe and Strzelecki Group's sediments.

On the basis of the new palynological data, a new well correlation has been made which exhibits the following:

Palaeocene sequence, the Childers Formation equivalent, is present along the north-eastern trend of Permit 109 and attenuates towards the west and over the Burong Fault Block. This sequence, however, is absent over the North Seaspray structure. Burong-1 has penetrated 50 feet of this sequence at around 4,050 feet (1,234 metres) K.B. The well has bottomed in the Strzelecki Group of Albian age (c. Paradoza zone).

Late Cretaceous sequence, the Golden Beach Formation equivalent, is present along the northeasterly trend of the coastline. This sequence, however, is absent over the Seaspray-Burong-North Seaspray Block and also on the Wellington Park structure. It is important to note that the study has confirmed the presence of the Late Cretaceous sequence in Carrs Creek-1 and Dutson Downs-1 wells, as predicted previously in our report. The study has also revealed that these wells have only penetrated the Early Cretaceous rocks at T.D.

DST tests over the Late Cretaceous (Golden Beach Formation) in Merriman-1 and Carrs Creek-1 at 5,070 feet (1,545 metres) and 4,600 feet (1,402 metres) respectively recovered water with Na Cl salinity of 15,850 ppm and 11,000 ppm, which indicates that the flushing probably has not affected the sediments of this sequence in the onshore area. This sequence was previously considered as Strzelecki Group by the operators.

The spore colours examined by the palynologist suggests that the Palaeocene rocks in the onshore area are immature to marginally mature for hydrocarbon generation, while the Late Cretaceous rocks are early mature to marginally mature. The results of the vitrinite reflectance measurements are broadly consistent with the results of the Rock-Eval Pyrolysis. However, lack of data below 5,000 feet (1,500 metres) in some of the wells and the availability of suitable samples has hampered the meaningful extrapolation of data, so that the results are not a true representation of the maturity profile in most of the wells.

The results of the extrapolation of the vitrinite reflectance have been plotted on the well correlation chart, which shows that the higher maturity gradients in the Merriman, North Seaspray and Wellington Park structures as compared to other locations may be related to the higher heat flows at these structures. The potential yield (i.e. $S_1 + S_2$) of these sediments has been calculated and indicates that the sediments of the Latrobe Group with an average of 14.63 kg hydrocarbon/tone rock have the highest source richness as compared to the Strzelecki Group with an average of 5.11 kg hydrocarbon/tone rock. The results of the potential yield for each formation have been calculated and are as follows:

Formation	(S₁ + S₂) range kg/tonne	(S₁ + S₂) average	kg/tonne
Latrobe Group	10.14-25.80	14.63	
Childers Formation (Palaeocene)	.15-21.82	6.77	
Golden Beach Formation (Late Cretaceous)	.76-18.45	9.60	
Strzelecki Group (Early Cretaceous)	.89-13.53	5.11	

All the sediments from Eocene to Early Cretaceous contain organic matter with the bulk composition of predominantly gas prone type III Kerogen. They consist largely of vitrinite with moderate proportions of inertinite and exinite. However, these gas prone sediments have the potential to generate liquid hydrocarbons in a significant quantity at optimum maturity.

Furthermore, light oil/condensate can be generated from the thermally labile exinite with VR=0.45% at around 2,300 metres subsea with the oil window (VR=0.7%) at around 3,600 metres subsea. Significant gas, however, can be generated from woody-herbaceous kerogen at as maturity around VR=0.6%, which corresponds approximately to 3,000 metres subsea. On the basis of these results, sediments of the Strzelecki Group are the only potential source rocks, which are mature below 2,300 metres subsea and can generate both gas and liquid hydrocarbons in the onshore area.

SAEED SHOGHI
CONSULTANT

27 March, 1991

APPENDIX 1

**PALYNOLOGY OF 23 SUBSURFACE SAMPLES,
ONSHORE GIPPSLAND BASIN
BY ROGER MORGAN**

PALYNOLOGY OF 23 SUBSURFACE SAMPLES,

GIPPSLAND BASIN

BY

ROGER MORGAN
BOX 161
MAITLAND, SOUTH AUSTRALIA, 5573

PHONE: (088) 322795
FAX: (088) 322798
REF: SD.GIP.CLUFFGIP

FOR CLUFF RESOURCES

FEBRUARY, 1991.

PALYNOLOGY OF 23 SUBSURFACE SAMPLES,
GIPPSLAND BASIN.

BY

ROGER MORGAN

<u>CONTENTS</u>	<u>PAGE</u>
I SUMMARY	3
II INTRODUCTION	5
III PALYNOSTRATIGRAPHY	7
IV CONCLUSIONS	15
V REFERENCES	17

I SUMMARY

The samples studied yielded the following zonal assignments.

Burong-1 :

4100 ft. (cutts) : mixed L. balmei zone (Paleocene) with minor late Cretaceous

4120 ft. (cutts) : C. paradoxa zone (late Albian)

Carrs Creek-1 :

3860 ft. (cutts), 4470 ft. (cutts) : mixed Tertiary, mostly Eocene

4570 ft. (cutts) : C. triplex zone (Coniacian-Turonian)

Dutson Downs-1 :

4590 ft. (cutts) : apparently L. balmei zone (Paleocene)

4740 ft. (cutts), 5360 ft. (cutts) : T. longus (Maastrichtian) to N. senectus (Campanian)

6020 ft. (cutts) : mixed C. triplex (Coniacian-Turonian) and C. striatus (early Albian)

Merriman-1 :

4708-14 ft. (CORE) : T. longus zone (Maastrichtian)

5057-81 ft. (CORE), 5740 ft. (cutts), 5950 ft. (cutts) : apparently all C. triplex zone (Coniacian-Turonian)

North Seaspray-1 :

3650 ft. (cutts) : C. paradoxa zone (late Albian)

3840 ft. (cutts), 4060 ft. (cutts) : C. striatus zone

(early Albian)

Seaspray-1 :

4590 ft. (cutts) : T. longus zone (Maastrichtian)

4790 ft. (cutts), 4879 ft. (cutts) : C. paradoxa zone (late Albian)

Wellington Park-1 :

3719-39 ft. (CORE) : L. balmei zone (Paleocene)

3816-19 ft. (CORE), 4550 ft. (cutts) : apparently all C.
paradoxa zone (late Albian)

7380 ft. (CORE) : indeterminate mid Jurassic to mid
Cretaceous

These data provide palynological confirmation for the Golden Beach Formation in Carrs Creek-1, Dutson Downs-1, and Merriman-1 and suggest its absence from the other wells. Heavy caving of the Tertiary in the cuttings samples could have masked older assemblages, however.

II INTRODUCTION

Babek Vazhebdedh of Cluff Resources submitted 23 samples for palynology from 7 Gippsland Basin wells. The study was aimed to test for the presence of late Cretaceous strata equivalent to the Golden Beach Formation, as discussed by Lowry (1987). Raw data is presented in Appendix I.

The published palynostratigraphic framework for the Cretaceous of Australia is most recently reviewed by Helby, Morgan and Partridge (1987). Dinoflagellates had been only rarely recorded from the Cretaceous of the Gippsland Basin, although Marshall (1988) provided taxonomic study of some Santonian dinoflagellates. In unpublished work, Marshall (1987a) describes dinoflagellates from new cuttings samples in Pisces-1, Marshall (1987b) describes taxonomy and some stratigraphy of Campanian dinoflagellates and in (1987c) describes some Santonian algal cysts. These all provide clues to the Coniacian-Turonian dinoflagellate sequence, but none provides the basis for a working zonation. The zonal scheme of Helby, Morgan and Partridge is shown in figure 1.

In the Tertiary, the Gippsland zonal scheme was most recently published by Partridge (1976), but the scheme is essentially similar to that for New Zealand for which substantial new data is available in Wilson (1988). Significant new Gippsland data is available in unpublished and privately circulated material, Harris (1985), Morgan (1988) and Marshall and Partridge (1988). The zonal framework of Partridge (1976) is shown in fig.1.

Organic maturity data was generated in the form of the Spore Colour Index. The oil and gas windows follow the general consensus of geochemical literature. The oil window corresponds to spore colours of light-mid brown (2.7) to

dark brown (3.6). This would correspond to Vitrinite Reflectance values of 0.6% to 1.3%. However, factors such as detailed kerogen type, basin type, basin history and heating curves all affect precise interpretation, and analytical machine-based maturity parameters are probably more reliable.

AGE	SPORE - POLLEN ZONES		DINOFLAGELLATE ZONES
Early Tertiary	Early Oligocene	<i>P. tuberculatus</i>	
	Late Eocene	upper <i>N. asperus</i>	<i>P. comatum</i>
		middle <i>N. asperus</i>	<i>V. extensa</i>
	Middle Eocene	lower <i>N. asperus</i>	<i>D. heterophycata</i>
			<i>W. echinosuturata</i>
		<i>P. asperopolus</i>	<i>W. edwardsii</i>
			<i>W. thompsonae</i>
		upper <i>M. diversus</i>	<i>W. ornata</i>
			<i>W. walpawaensis</i>
	Early Eocene	middle <i>M. diversus</i>	
		lower <i>M. diversus</i>	<i>W. hyperacantha</i>
Late Cretaceous	Paleocene	upper <i>L. balmei</i>	<i>A. homomorpha</i>
		lower <i>L. balmei</i>	<i>E. crassitabulata</i>
	Maastrichtian		<i>T. evittii</i>
		<i>T. longus</i>	<i>M. druggii</i>
	Campanian	<i>T. lilliei</i>	<i>I. korjonense</i>
		<i>N. senectus</i>	<i>X. australis</i>
	Santonian	<i>T. pachyexinus</i>	<i>N. aceras</i>
			<i>I. cretaceum</i>
	Coniacian		<i>O. porifera</i>
	Turonian	<i>C. triplex</i>	<i>C. striatoconus</i>
			<i>P. infusoriooides</i>
Early Cretaceous	Cenomanian	<i>A. distocarinatus</i>	
	Albian	<i>P. pannosus</i>	
		upper <i>C. paradoxa</i>	
		lower <i>C. paradoxa</i>	
	Aptian	<i>C. striatus</i>	
		upper <i>C. hughesi</i>	
		lower <i>C. hughesi</i>	
	Barremian		
	Hauterivian	<i>F. wonthaggiensis</i>	
	Valanginian	upper <i>C. australiensis</i>	
	Berriasian	lower <i>C. australiensis</i>	
Juras	Tithonian	<i>R. watherooensis</i>	

FIGURE 1

ZONATION FRAMEWORK

Ma




Figure 2 — Time-space diagram generalised for Cretaceous to Eocene in the offshore Gippsland Basin.

III PALYNOSTRATIGRAPHY

A. BURONG-1

- 4100 ft. (cutts) : mixed L. balmei zone and younger with minor late Cretaceous

The age is uncertain as rare Eocene restricted taxa (P. pachypolus and M. tenuis) occur mixed with Paleocene and older taxa (L. balmei, G. rudata, T. verrucosus) and very rare Maastrichtian and older taxa (T. confessus). H. harrisii is dominant with frequent G. rudata and Nothofagidites spp. The most obvious interpretation is a Paleocene balmei zone assignment with minor reworked Cretaceous. However, a Maastrichtian upper longus zone assignment is also possible. Nothing older was seen.

Environments appear to be non-marine with abundant and diverse spores and pollen and no marine indicators.

Light brown spore colours suggest marginal maturity for oil generation.

- 4120 ft. (cutts) : C. paradoxa Zone

This assemblage is relatively clean, dominated by O. wellmannii and assigned on the presence of C. paradoxa and the associated spore dominated floras including Aequitriradites spp., C. australiensis, F. dailyi and T. trioreticulosus. Rare Triassic reworking was seen.

Environments are non-marine fluvial on account of the abundant and diverse spore dominated assemblage and absence of marine indicators.

Light to mid brown spore colours suggest early maturity

for oil generation.

B. CARRS CREEK-1

- 3860 ft. (cutts), 4470 ft. (cutts) : mixed Tertiary,
mostly Eocene

These assemblages are dominated by H. harrisii,
Nothofagidites spp. and Proteacidites spp. Eocene
indicators include M. diversus, M. tenuis, P.
pachypolus and common N. deminatus. Nothing older
than Eocene was seen, but heavy caving may be masking
something older.

- 4570 ft. (cutts) : C. triplex Zone

This assemblage is totally dominated by small
Dilwynites and contains frequent P. mawsonii. More
important, it contains the distinctive algal
Rimosicysta spp, which are P. mawsonii zone restricted.

Non-marine environments are indicated by the absence of
saline indicators, but lacustrine influence is
suggested by the algal assemblage.

Light brown spore colours suggest marginal maturity for
oil.

C. DUTSON DOWNS-1

- 4590 ft. (cutts) : apparently L. balmei zone

Assignment is made on L. balmei without older
indicators. H. harrisii and N. emarcidus are common.

Non-marine environments are suggested by the common

cuticle and diverse spores and pollen without marine indicators.

Yellow to light brown spore colours indicate immaturity for hydrocarbons.

- 4740 ft. (cutts)-5360 ft. (cutts) : apparently T. longus to N. senectus zones

These samples are almost identical with that above apart from the presence of rare Late Cretaceous forms including T. sabulosus (longus to senectus restricted). T. longus (longus restricted) and T. confessus (longus to pachyexinus restricted). It is interpreted as upper late Cretaceous with heavy Tertiary caving, but caving is so heavy that it could be older and masked.

Non-marine environments are indicated by the lack of marine indicators and the abundant cuticle and spore-pollen.

Light brown spore colours suggest marginal maturity for oil.

- 6020 ft. (cutts) : mixed C. triplex zone with C. striatus zone

The early Cretaceous striatus zone is indicated by the association of C. striatus with P. notensis without C. paradoxa. Other spores supporting the Early Cretaceous age include C. holodictyus, Aequitriradites spp., F. wonthaggiensis and Triporoletes spp. The Turonian-Coniacian triplex zone is indicated by the algal Rimosicysta and frequent P. mawsonii but are considered caved. The triplex zone must therefore occur above this point in the well. Dominant however,

is the Paleocene balmei assemblage, caving very heavily.

Non-marine striatus environments are non-marine.

Lacustrine triplex environments are suggested by the associations described above.

Light to mid brown spore colours suggest early maturity in the striatus zone assemblages.

D. MERRIMAN-1

- 4708-14 ft. (CORE) : upper T. longus zone

This lean and cuticle dominated assemblage is assigned at the base on oldest S. punctatus and at the top on youngest O. sentosa and frequent G. rudata, and the absence of L. balmei. Frequent forms include G. rudata and S. punctatus.

Non-marine possibly lacustrine environments are indicated by the very rare nondescript dinoflagellates in the cuticle and spore-pollen dominated assemblage.

Yellow to light brown spore colours suggest early marginal maturity for oil.

- 5057-81 ft. (CORE), 5740 ft. (cutts), 5950 ft. (cutts) : apparently all C. triplex zone

Assignment of the core is straightforward on youngest C. "pileosus" and oldest P. mawsonii. Common forms include P. mawsonii and M. antarcticus. The two cuttings samples beneath contain heavy Paleocene balmei caving, but also contain triplex zone indicators (common Dilwynites and algal Rimosicysta spp) and

nothing older. They are therefore assigned to the triplex zone.

Non-marine environments with some lacustrine influence are indicated by the algal acritarchs, diverse pollen and spores and abundant cuticle.

Yellow to light brown spore colours indicate early marginal maturity for oil.

E. NORTH SEASPRAY-1

- 3650 ft. (cutts) : C. paradoxa zone

Although the assemblage is dominated by Paleocene L. balmei zone and Eocene diversus zone taxa, the rare presence of Aequitriradites spp., C. paradoxa, C. striatus, P. linearis and T. reticulatus indicate the paradoxa zone.

Non-marine environments are indicated by the abundant cuticle, dominant spore-pollen, and absence of marine indicators.

Light brown spore colours indicate marginal maturity for hydrocarbons.

- 3840 ft. (cutts), 4060 ft. (cutts) : C. striatus zone

These sample are almost identical with that above, but amongst the rare elements, markers for the paradoxa zone cannot be found. Oldest C. striatus therefore indicates the older striatus zone, although its base could be caved.

Non-marine environments and marginal maturity for oil

are indicated as above.

F. SEASPRAY-1

- 4590 ft. (cutts) : probably T. longus zone

Although Paleocene L. balmei taxa dominate the sample, the rare occurrence of Cretaceous taxa such as N. senectus (longus to senectus restricted), T. verrucosus (balmei to longus restricted) and T. sectilis (longus to lillei restricted) indicate the longus zone assignment.

Non-marine environments are indicated by the absence of marine indicators and the diverse pollen and spores.

Yellow to light brown spore colours indicate early marginal maturity for oil.

- 4790 ft. (cutts), 4879 ft. (cutts) C. paradoxa zone

Although again Paeocene caving is dominant, rare Aequitirradites, C. paradoxa, C. striatus, Foraminispora spp. and Triporoletes spp. indicate the paradoxa zone.

Non-marine environments are indicated by diverse spores and pollen. The single spiny acritarch at 4879 ft. is considered caved.

Light to mid brown spore colours indicate early maturity for oil.

G. WELLINGTON PARK-1

- 3719-39 ft. (CORE) : L. balmei zone

Assignment is clearly indicated at the top by youngest L. balmei and T. verrucosus and at the base by oldest S. punctatus and T. verrucosus without older indicators. Common forms are P. mawsonii and Proteacidites spp. A single T. waiparaensis is considered reworked.

Non-marine environments are indicated by the diverse pollen and spores and absence of marine indicators.

Yellow spore colours indicate immaturity for oil generation.

- 3816-19 ft. (CORE), 4550 ft. (cutts) : apparently all C. paradoxa zone

Assignment of the core sample is straightforward on youngest P. notensis (and the other Early Cretaceous associates) and oldest C. paradoxa. Common forms include Cyathidites and Falcisporites. Permian and Triassic reworking were seen. The cuttings at 4550 ft. contain the same assemblage but could clearly be caved into something slightly older.

Non-marine environments are indicated on the absence of saline indicators and the diverse pollen and spores.

Light to mid brown spore colours indicate early maturity for oil.

- 7380 ft. (CORE) : Jurassic-Cretaceous : indeterminate

This sample is very lean of palynomorphs and all are longranging taxa. The presence of C. dampieri, C. penolaensis and R. nodosus indicates a Middle Jurassic

to mid Cretaceous age range, but more precision is not possible.

Non-marine environments are suggested, but too few palynomorphs have been seen to be confident.

Mid to dark brown spore colours indicate peak maturity for oil generation.

IV CONCLUSIONS

Clearly the current study was directed use palynology to test the log based already mapped distribution of the Golden Beach Formation. The Gippsland sequence along the northern margin usually comprises the early Cretaceous hughesi to paradoxa Strzelecki Group, a pannosus-distocarinatus unconformity corresponding to Southern Ocean breakup, an early Late Cretaceous triplex Golden Beach Formation, a pachyexinus-senectus unconformity corresponding to Tasman Sea breakup, and the lillei to asperus Latrobe Group. This is discussed in more detail in Lowry (1987) and summarized here in figure 2.

In all the cuttings samples, the younger Latrobe Group caves heavily and may confuse interpretation. Nevertheless, most wells are straight-forward.

In Burong-1, longus Latrobe Group appears to directly overlie paradoxa Strzelecki Group on an unconformity in the gap 4100 ft. to 4120 ft. The Golden Beach Formation appears to be absent.

In Carrs Creek-1, only cuttings are available, and these may mask the true situation. On the palynology, Eocene Latrobe Group appears to directly overlie triplex Golden Beach Formation, with the unconformity in the gap 4470 ft. to 4570 ft.

In Dutson Downs-1, again only cuttings are available and appear to have masked assemblages, making them appear to be deeper. The sequence appears to comprise balmei to possibly senectus Latrobe group (4590 to 5360 ft.), triplex Golden Beach Formation (somewhere in the gap 5360 to 6020 ft.) and Strzelecki Group (6020 ft.). Logs suggest Golden Beach Formation at 1442-1781m (4731-5848 ft.) (Lowry 1987)

and may well be right.

In Merriman-1, longus Latrobe Group overlies triplex Golden Beach Formation with the unconformity in the gap 4714 ft. to 5057 ft. The well appears to still be in Golden Beach Formation at 5950 ft., but these cuttings could be masking something older.

In North Seaspray-1, Strzelecki Group spans the sampled interval, although balmei-diversus Latrobe Group caving is heavy.

In Seaspray-1, longus Latrobe Group directly overlies paradoxa Strzelecki Group with the unconformity in the gap 4590 ft. to 4790 ft. The Golden Beach Formation is missing.

In Wellington Park-1, balmei to possibly longus Latrobe Group directly overlies paradoxa Strzelecki Group with the unconformity in the gap 3739 to 3816 ft. The Golden Beach Formation is missing.

V

REFERENCES

Harris, W.K. (1985) Middle to Late Eocene Depositional Cycles and Dinoflagellate Zones in Southern Australia Spec. Publ., S. Aust. Dept. Mines and Energy 5 : 133-144

Helby, R.J., Morgan, R.P., and Partridge A.D., (1987) A palynological Zonation of the Australian Mesozoic Australas. Assoc. Palaeont., Mem. 4

Lowry, D.C.(1987) A new play in the Gippsland Basin Apea J. 27 (1) 164 - 172.

Marshall, N. (1987a) Palynological analysis of Pisces-1, Gippsland Basin, southeastern Australia Earth Resources Foundation, Sydney University unpubl. rpt. 1987/1

Marshall, N. (1987b) Campanian dinoflagellates from southeastern Australia Earth Resources Foundation, Sydney University, unpubl. rpt. 1987/3

Marshall, N. (1987c) An unusual assemblage of algal cysts from the Turonian - early Santonian of the Gippsland Basin, southeastern Australia Earth Resources Foundation, Sydney University unpubl. rpt. 1987/5

Marshall, N.G. (1988) A Santonian dinoflagellate assemblage from the Gippsland Basin, southeastern Australia Australas. Assoc. Palaentols. Mem. 5, 195-213

Marshall, N.G. and Partridge A.D. (1988) The Eocene acritarch Tritonites gen. nov. and the age of the Marlin Channel, Gippsland Basin, southeastern Australia Australas. Assoc. Palaentols. Mem. 5, 239-257

Partridge A.D. (1976) The Geological Expression of
Eustacy in the Early Tertiary of the Gippsland Basin
Aust. Pet. Explor. Assoc. J., 16 : 73-79

PALYNOLOGICAL DATA OF 7 WELLS incl. BURONG #1, MERRIMAN #1

ROGER MORGAN : PALYNOLOGY CONSULTANT
BOX 161, MAITLAND, SOUTH AUSTRALIA, 5573
PHONE (088) 322795 FAX (088) 322798

C L I E N T: Cluff Resources

W E L L: 7 inc. Burong, Carr's Creek, Merriman, Nth Seaspray

F I E L D / A R E A: Gippsland Basin (Northen Margin)

A N A L Y S T: Roger Morgan

D A T E : February '91

N O T E S: all sample depths are in metres

RANGE CHART OF GRAPHIC ABUNDANCES BY GROUP: DINOS and S/POLLEN

Key to Symbols

- = Very Rare
- = Rare
- = Few
- = Common
- = Abundant
- ? = Questionably Present
- . = Not Present

= Abundant

? = Questionably Present

. = Not Present

1	DEFLANDREA SPP
2	MICHYRTRIDIUM
3	OPERCULODINIUM SPP
4	RIMOSICYSTA ASPERA
5	RIMOSICYSTA EVERSA
6	SCHIZOSPORIS
7	SUBTILISPHAERA
8	AEQUITIRADITES HISPIUS
9	AEQUITIRADITES SPINULOSUS
10	AEQUITIRADITES TILCHAENESIS
11	AEQUITIRADITES VERRUCOSUS
12	AMOSOPOLLIS CRUICIFORMIS
13	ANNULISPORITES
14	APPENDICISPORITES DISTOCARINATUS
15	ARAUCARIACITES AUSTRALIS
16	AUSTRALOPOLLIS OBSCURUS
17	BACULATISPORITES
18	BALMEISPORITES GLENELGENSIS
19	BALMEISPORITES HOLODICTYUS
20	BALMEISPORITES TRIDICTYUS
21	BANKSIEACIDITES ARCUATUS
22	CALLIALASPORITES DAMPIERI
23	CAMEROZONOSPORITES AMBIGENS
24	CAMEROZONOSPORITES SP
25	CERATOSPORITES EQUALIS
26	CICATRICOSISPORITES AUSTRALIENSIS
27	CICATRICOSISPORITES CUNEIFORMIS
28	CICATRICOSISPORITES HUGHESI
29	CLAVIFERA TRIPLEX
30	CONCAVISSIMISPORITES PENOLAENSIS
31	COPTOSPORA PARADOXA
32	COPTOSPORA PILEOSUS
33	COPTOSPORA SP

BURONG-1

4100 CUTT

4120 CUTT

CARRS CREEK-1

3850-60 CUTT

4460-70 CUTT

4560-70 CUTT

DUTSN DOWNS-1

4580-90 CUTT

4730-40 CUTT

5350-60 CUTT

5010-20 CUTT

MERRIMAN-1

4708-14 CORE

5075-81 CORE

5730-40 CUTT

5940-50 CUTT

NTH SEASPRY-1

3640-50 CUTT

3830-40 CUTT

4050-60 CUTT

SEASPRAY-1

4580-90 CUTTS

4780-90 CUTTS

4872-79 CORE

WELLNGTN PK-1

3719-39 CORE

3816-19 CORE

4540-50 CUTT

7379-80 CORE

67	LKLKISPORITES SCABERIS
68	LAEVIGATOSPORITES OVATUS
69	LATROBOSPORITES AMPLUS
70	LATROBOSPORITES CRASSUS
71	LATROBOSPORITES OHAIENSIS
72	LEPTOLEPIDITES VERRUCATUS
73	LILIACIDITES PERORETICULATUS
74	LYCOPODIACIDITES ASPERATUS
75	LYGISTEPOLLENITES BALMEI
76	LYGISTEPOLLENITES ELLIPTICUS
77	LYGISTEPOLLENITES FLORINII
78	MALVACIPOLLIS DIVERSUS
79	MALVACIPOLLIS SUBTILIS
80	MATONISPORITES COOKSONIAE
81	MICROCACHRYDITES ANTARCTICUS
82	MYRTACEIDITES TENUIS
83	NEORAISTRICKIA
84	NOTHOFAGIDITES
85	NOTHOFAGIDITES BRACHYSPINULOSUS
86	NOTHOFAGIDITES DEMINUTUS
87	NOTHOFAGIDITES EMARCIDUS / HETERUS
88	NOTHOFAGIDITES ENDURUS
89	NOTHOFAGIDITES FALCATUS
90	NOTHOFAGIDITES FLEMINGII
91	NOTHOFAGIDITES SENECTUS
92	ORNAMENTIFERA SENTOSA
93	OSMUDACIDITES WELLMANII
94	PERIPOROPOLLENITES POLYORATUS
95	PERIPOROPOLLENITES SPP
96	PERIMOPOLLENITES VESICUS
97	PEROTRILETES LINEARIS
98	PEROTRILETES MAJUS
99	PHIMOPOLLENITES PANNOUS

BURONG-1

4100 CUTT

4120 CUTT

CARRS CREEK-1

3850-60 CUTT

4460-70 CUTT

4560-70 CUTT

DUTSN DOWNS-

4580-90 GUTT

4730-40 CUTT

5350-60 CUTT

6010=20 CU

MERRITMAN-1

MERRIMAN-1
4308-14 CORP

4708-14 CORE
5035-81 CORE

5075-81 CURE
5330-49 CUT

5730-40 CUTT
5840-50 CUTT

3940-30 COT
NTU SEARCHED

NIA SEASPRY-
3440 EO CUTT

3640-30 CUT
3830-40 CUT

3830-40 CU

4050-60 COT
REARRAY 1

SEASPRAY-1

4580-90 CUT
4380-80 CUT

4780-90 CUT

4872-79 CORE
HELIANTHUS RM

WELLNGTN PK-1
78148 00 0000


3719-39 CORE


3816-19 CORE
1540 E9 QUETZ

4540-50 COT

APPENDIX 2

**REPORT ON SOURCE ROCK STUDY ON 36 SAMPLES
FROM ONSHORE GIPPSLAND
BY AMDEL CORE SERVICES**





6th March 1991

Cluff Resources Pacific Ltd
PO Box 908
NORTH SYDNEY NSW 2060

Attention: Babak Vahebzadeh

REPORT: 009/732

CLIENT REFERENCE: Letter 8/1/91

MATERIAL: Cuttings and Core

LOCALITY: Onshore Gippsland Basin

WORK REQUIRED: TOC, Rock-Eval Pyrolysis and Vitrinite Reflectance Determinations

Please direct technical enquiries regarding this work to the signatory below under whose supervision the work was carried out.

A handwritten signature in black ink, appearing to read "Brian L. Watson".

BRIAN L WATSON
Laboratory Supervisor
on behalf of Amdel Core Services Pty Ltd

Amdel Core Services Pty Limited shall not be liable or responsible for any loss, cost, damages or expenses incurred by the client, or any other person or company, resulting from any information or interpretation given in this report. In no case shall Amdel Core Services Pty Ltd be responsible for consequential damages including, but not limited to, lost profits, damages for failure to meet deadlines and lost production arising from this report.

1. INTRODUCTION

Thirty six core and cuttings samples from the onshore Gippsland Basin were received for geochemical analyses to determine the source richness, source maturity and source quality of these sediments.

This report is a formal presentation of the results of this work which was reported by facsimile on the 8th and 21st of February 1991 and includes an interpretation of the complete data set along with graphical displays of the data.

2. ANALYTICAL PROCEDURES

Analytical procedures used in this study are presented in Appendix 2.

3. RESULTS

TOC and Rock-Eval pyrolysis data is presented in Table 1. Figure 1 is a plot of Rock-Eval Hydrogen index versus Tmax illustrating kerogen Type and maturity. Table 2 is a summary of the Vitrinite Reflectance determinations. Histogram plots of these determinations are presented in Appendix 1. Plots of Vitrinite Reflectance versus depth are included as Figures 2-7.

4. INTERPRETATION

4.1 Maturity

Vitrinite Reflectance measurements (Table 2) indicate that the sediments examined in this study, are for the most part, immature for the generation of hydrocarbons. Maturities indicated by Rock-Eval pyrolysis Tmax values are broadly consistent with these indications.

The amount of Vitrinite reflectance data available for each well is of course limited by the availability of suitable samples. However, it should be noted that extrapolation of this data is limited in some cases by the spread of values with depth. The lack of data below 5,000 feet depth in Burong-1, Seaspray-1 and North Seaspray presents some difficulties in extrapolating this data to depths in excess of 8,000 feet.

The validity of the apparent change in slope in the Wellington Park-1 Reflectance versus depth plot should be further investigated. The data available indicates that the rate of change of thermal maturity changes with depth as indicated. Curved Reflectance versus depth plots are common, particularly when approaching sedimentary basement or intrusives. Vitrinite Reflectance determinations of further sediments below 6000 feet depth would be advisable to substantiate the indications of the present data set.

Extrapolation of the available Vitrinite Reflectance data indicates that the sediments intersected in these wells will be sufficiently mature for the generation of light oil/condensates from the thermally labile exinites (resinite and bituminite) below the following depths (VR threshold = 0.45%):

<u>Well</u>	<u>Approximate Depth (VR = 0.45%)</u>
Burong - 1	15,000 ft
Dutson Downs	8,000 ft
Merriman - 1	6,000 ft
North Seaspray - 1	6,500 ft
Seaspray - 1	9,500 ft
Wellington Park - 1	6,300 ft

Significant gas generation from woody - herbaceous kerogen commences at maturities corresponding to Vitrinite Reflectance values of approximately 0.6%. The available data indicates that sediments in these locations are sufficiently mature for significant gas generation below approximately:

<u>Well</u>	<u>Approximate Depth (VR - 0.6%)</u>
Burong - 1	Insufficient data
Dutson Downs - 1	12,000 - 13,000 ft
Merriman - 1	9,000 ft
North Seaspray - 1	11,000 ft
Seaspray-1	20,000 ft
Wellington Park - 1	7,300 ft

The generation of oil from exinites other than resinite and bituminite commences of maturities corresponding to a Vitrinite Reflectance of 0.7%. Extrapolation to this maturity is not possible in the majority of the wells examined in this study, however, the available data suggests that sediments intersected in the following locations are sufficiently mature for oil generation below approximately:

<u>Well</u>	<u>Approximate Depth (VR = 0.7%)</u>
Merriman - 1	13,000 ft
North Seaspray - 1	13,000 - 15,000 ft
Wellington Park - 1	7,600 ft

Vitrinite Reflectance versus depth plots (Figures 2 - 7) indicate higher maturity gradients at the following locations: Merriman - 1, North Seaspray - 1 and Wellington Park - 1 as compared to maturity gradients at the other locations examined (Burong - 1, Dutson Downs - 1 and Seaspray - 1). Higher maturity gradients indicate higher heat flows and should be related to the geology of the region for maximum benefit. Understanding these heat flows may lead to the identification of a region where sediments are optimally mature for hydrocarbon generation at shallower depths.

Rock-Eval production indices of greater than 0.2 indicate that the following samples contain migrated hydrocarbons:

Dutson Downs - 1	5360 feet depth
Wellington Park - 1	3650 - 7376 feet depth

4.2 Source Richness

Organic richness ranges from poor to excellent in the sediments studied (TOC = 0.18 - 49.60%; Table 1). TOC values are typically greater than 1% and are less than 1% in only eight of the thirty-six samples analysed. The absence of coal samples from Burong - 1 and Seaspray - 1 results in a data set which suggests a lower organic richness for sediments in these wells. However, these observations may be related to sampling or the availability of sample and should be further evaluated.

Source Richness for the generation of hydrocarbons also ranges from poor to excellent ($S_1 + S_2 = 0.15 - 50.48$ kg of hydrocarbons/tonne), but is generally fair to excellent (>2 kg of hydrocarbons/tonne).

4.3 Kerogen Type and Source Richness

Hydrogen Index and Tmax data (Table 1; Figure 1) indicates that these sediments contain organic matter with the bulk composition of predominantly gas prone Type III kerogen. Visual examination of the organic matter in these samples shows that whilst they consist largely of vitrinite they contain moderate proportions of both inertinite and exinite (5 - 15% of organic matter).

This data suggests that whilst these sediments are likely to be largely gas prone they do have the potential to generate significant quantities of liquid hydrocarbons on maturity.

TABLE 1

Page No 1

AMDEL CORE SERVICES

Rock-Eval Pyrolysis

08/02/91

Client: CLUFF RESOURCES

Basin: GIPPSLAND (ONSHORE)

Depth (ft)	T Max	S1	S2	S3	S1+S2	PI	S2/S3	PC	TOC	HI	OI
NORTH SEASPRAY-1											
1910	380	0.42	14.89	18.61	15.31	0.03	0.80	1.27	49.60	30	37
2100	385	0.83	7.66	3.28	8.49	0.10	2.33	0.70	15.60	49	21
3504	382	0.48	12.78	7.30	13.26	0.04	1.75	1.10	29.90	42	24
3690	380	0.16	3.33	3.53	3.49	0.05	0.94	0.27	11.40	29	30
3800	384	0.07	0.44	1.76	0.51	0.14	0.25	0.03	3.66	12	48
4000	401	0.06	1.70	1.18	1.76	0.03	1.44	0.14	3.01	56	39
5000	221	0.02	0.21	0.58	0.23	0.09	0.01	0.00	0.61	35	95
MERRIMAN-1											
2210									0.38		
2510	403	0.39	17.87	7.00	18.26	0.02	2.55	1.52	40.10	44	17
4350	412	0.46	10.23	2.31	10.69	0.04	4.42	0.89	11.10	92	20
4720	372	0.01	0.73	6.75	0.74	0.01	0.10	0.06	5.45	13	123
5081	339	0.01	0.75	2.36	0.76	0.01	0.31	0.06	1.45	51	162
5960	381	0.38	4.15	1.89	4.53	0.08	2.19	0.37	2.78	149	67
DUTSON DOWNS-1											
2420	380	1.03	21.01	8.17	22.04	0.12	2.35	2.07	14.60	143	61
3070	385	3.30	47.18	9.22	50.48	0.07	5.11	4.20	25.40	185	36
4210	389	0.22	4.68	1.39	4.90	0.04	3.36	0.40	3.81	122	36
4660	413	1.29	20.53	8.58	21.82	0.06	2.39	1.81	10.60	193	80
4820	390	2.68	32.46	8.98	35.14	0.08	3.61	2.92	24.20	134	37
5360	395	0.40	1.37	2.02	1.77	0.23	0.67	0.14	7.66	17	26
6110	372	0.46	4.66	1.87	5.12	0.09	2.49	0.42	1.25	372	149
BURONG-1											
2100									0.26		
2670	389	0.88	21.45	2.87	22.33	0.04	7.47	1.86	8.69	246	33
3950	406	0.14	1.82	0.42	1.96	0.07	4.33	0.16	1.33	136	31
4120	376	0.00	0.15	0.40	0.15	0.00	0.37	0.01	0.40	37	100
WELLINGTON PARK-1											
1180									0.18		
1880									0.25		
2100									0.29		
3310	230	1.13	4.92	7.80	6.05	0.19	0.63	0.50	46.70	10	16
3650	283	1.35	3.98	5.33	5.33	0.25	0.74	0.44	24.40	16	21
3830	283	2.92	10.79	6.28	13.71	0.21	1.71	1.14	31.20	34	20
7376	283	0.63	2.59	0.56	3.22	0.20	4.62	0.26	14.80	17	3
SEASPRAY-1											
4466	383	0.51	11.92	2.96	12.43	0.04	4.03	1.21	12.60	95	23
4730	380	0.21	5.02	2.43	5.23	0.04	2.07	0.31	7.49	67	32
4885	381	0.52	2.99	0.12	3.51	0.15	2.49	0.29	0.59	51	20
5000	257	0.13	0.76	0.21	0.89	0.15	3.61	0.07	0.62	122	34

KEY TO ROCK-EVAL PYROLYSIS DATA SHEET

	<u>PARAMETER</u>	<u>SPECIFICITY</u>
T max	position of S ₂ peak in temperature program (°C)	Maturity/Kerogen type
S ₁	kg hydrocarbons (extractable)/tonne rock	Kerogen type/Maturity/Migrated oil
S ₂	kg hydrocarbons (kerogen pyrolysate)/tonne rock	Kerogen type/Maturity
S ₃	kg CO ₂ (organic)/tonne rock	Kerogen type/Maturity *
S ₁ + S ₂	Potential Yield	Organic richness/Kerogen type
PI	Production Index (S ₁ /S ₁ + S ₂)	Maturity/Migrated Oil
PC	Pyrolysable Carbon (wt. percent)	Organic richness/Kerogen type/Maturity
TOC	Total Organic Carbon (wt. percent)	Organic richness
HI	Hydrogen Index (mg h'c (S ₂)/g TOC)	Kerogen type/Maturity
OI	Oxygen Index (mg CO ₂ (S ₃)/g TOC)	Kerogen type/Maturity *

*Also subject to interference by CO₂ from decomposition of carbonate minerals.

TABLE 2: SUMMARY OF VITRINITE REFLECTANCE MEASUREMENTS,
ON SHORE GIPPSLAND BASIN

Depth (ft)	Mean Maximum Reflectance %	Standard Deviation	Range	Number of Determinations
BURONG - 1				
2100	0.31 (0.25)	0.00	0.30-0.31	2
2670	0.26	0.03	0.19-0.33	27
3950	0.30	0.04	0.22-0.38	23
4120	0.30	0.05	0.22-0.41	13
DUTSON DOWNS - 1				
2420	0.32	0.03	0.25-0.37	16
3070	0.28	0.03	0.22-0.34	29
4210	0.35	0.04	0.28-0.41	21
4660	0.35	0.06	0.23-0.46	16
4820	0.37	0.03	0.31-0.43	25
5360	0.35	0.05	0.28-0.47	20
6110	0.44	0.06	0.34-0.53	24
MERRIMAN - 1				
2210	0.28	0.06	0.21-0.34	5
2510	0.26	0.03	0.20-0.31	31
4350	0.32	0.04	0.24-0.40	26
4720	0.39	0.02	0.33-0.45	24
5081	0.44	0.03	0.41-0.47	2
5960	0.46	0.05	0.40-0.55	12
NORTH SEASPRAY - 1				
1910	0.28	0.01	0.25-0.31	22
2100	0.29	0.02	0.26-0.33	24
3310	0.34	0.04	0.28-0.44	24
3504	0.32	0.03	0.28-0.37	24
3690	0.32	0.04	0.25-0.40	24
3800	0.33	0.03	0.25-0.40	24
4000	0.39	0.05	0.31-0.45	10
5000	0.39	0.05	0.29-0.48	16
SEASPRAY - 1				
4466	0.36	0.04	0.29-0.44	22
4730	0.37	0.06	0.28-0.49	23
5000	0.37	0.02	0.33-0.41	16
WELLINGTON PARK - 1				
1880	0.36	0.07	0.27-0.50	9
3310	0.28	0.03	0.24-0.33	24
3650	0.34	0.03	0.28-0.40	16
3830	0.33	0.03	0.28-0.39	22
7376	0.63	0.07	0.46-0.77	30

FIGURE 1

HYDROGEN INDEX vs T max

Company : CLUFF RESOURCES
Basin : GIPPSLAND (ONSHORE)




FIGURE 2

VITRINITE REFLECTANCE VERSUS DEPTH
BURONG-1




FIGURE 3

VITRINITE REFLECTANCE VERSUS DEPTH
DUTSON DOWNS-1




FIGURE 4

VITRINITE REFLECTANCE VERSUS DEPTH
MERRIMAN-1




FIGURE 5

VITRINITE REFLECTANCE VERSUS DEPTH
NORTH SEASPRAY-1




FIGURE 6

VITRINITE REFLECTANCE VERSUS DEPTH
SEASPRAY-1





FIGURE 7

VITRINITE REFLECTANCE VERSUS DEPTH
WELLINGTON PARK-1



APPENDIX 1: HISTOGRAM

PLOTS OF VITRINITE REFLECTANCE

DETERMINATIONS, ONSHORE

GIPPSLAND BASIN

VITRINITE REFLECTANCE VALUES

Well Name: BURONG-1
Depth: 2100 ft

Sorted List

0.30
0.31

Number of values= 2

Mean of values 0.31
Standard Deviation 0.00

HISTOGRAM OF VALUES
Reflectance values multiplied by 100

30-32 **

VITRINITE REFLECTANCE VALUES

Well Name: BURONG-1
Depth: 2670 ft

Sorted List

0.19	0.25	0.28
0.21	0.25	0.29
0.22	0.26	0.29
0.22	0.26	0.30
0.23	0.26	0.31
0.24	0.26	0.31
0.25	0.27	0.33
0.25	0.27	
0.25	0.27	
0.25	0.28	

Number of values= 27

Mean of values 0.26
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

19-21	**
22-24	****
25-27	*****
28-30	*****
31-33	***

VITRINITE REFLECTANCE VALUES

Well Name: BURONG-1
Depth: 3950 ft

Sorted List

0.22	0.29	0.35
0.23	0.29	0.37
0.25	0.30	0.38
0.25	0.30	
0.26	0.30	
0.27	0.31	
0.28	0.32	
0.28	0.33	
0.28	0.33	
0.29	0.35	

Number of values= 23

Mean of values 0.30
Standard Deviation 0.04

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

22-24	**
25-27	***
28-30	*****
31-33	****
34-36	**
37-39	**

VITRINITE REFLECTANCE VALUES

Well Name: BURONG-1
Depth: 4120 ft

Sorted List

0.22 0.33
0.24 0.38
0.25 0.41
0.26
0.27
0.28
0.29
0.30
0.30
0.32

Number of values= 13

Mean of values 0.30
Standard Deviation 0.05

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

22-24 **
25-27 **
28-30 *****
31-33 **
34-36
37-39 *
40-42 *

VITRINITE REFLECTANCE VALUES

Well Name: DUTSON DOWNS-1
Depth: 2420 ft

Sorted List

0.25	0.33
0.28	0.34
0.29	0.35
0.29	0.35
0.29	0.36
0.29	0.37
0.31	
0.31	
0.32	
0.33	

Number of values= 16

Mean of values 0.32
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

25-27	*
28-30	*****
31-33	*****
34-36	****
37-39	*

VITRINITE REFLECTANCE VALUES

Well Name: DUTSON DOWNS-1
Depth: 3070 ft

Sorted List

0.22	0.27	0.29
0.24	0.27	0.29
0.25	0.27	0.30
0.25	0.27	0.30
0.25	0.27	0.31
0.25	0.28	0.31
0.26	0.29	0.32
0.26	0.29	0.33
0.26	0.29	0.34
0.27	0.29	

Number of values= 29

Mean of values 0.28
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

22-24	**
25-27	*****
28-30	*****
31-33	
34-36	*

VITRINITE REFLECTANCE VALUES

Well Name: DUTSON DOWNS-1
Depth: 4210 ft

Sorted List

0.28	0.35	0.41
0.29	0.35	
0.29	0.36	
0.31	0.37	
0.32	0.37	
0.33	0.38	
0.33	0.38	
0.33	0.38	
0.33	0.39	
0.34	0.40	

Number of values= 21

Mean of values 0.35
Standard Deviation 0.04

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

28-30	***
31-33	*****
34-36	****
37-39	*****
40-42	**

VITRINITE REFLECTANCE VALUES

Well Name: DUTSON DOWNS-1
Depth: 4660 ft

Sorted List

0.23	0.37
0.28	0.37
0.28	0.41
0.29	0.43
0.33	0.44
0.33	0.46
0.33	
0.34	
0.35	
0.36	

Number of values= 16

Mean of values 0.35
Standard Deviation 0.06

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

23-25	*
26-28	
29-31	***
32-34	****
35-37	*****
38-40	
41-43	**
44-46	**

VITRINITE REFLECTANCE VALUES

Well Name: DUTSON DOWNS-1
Depth: 4820 ft

Sorted List

0.31	0.37	0.40
0.33	0.37	0.40
0.33	0.37	0.41
0.33	0.37	0.41
0.34	0.37	0.43
0.34	0.37	
0.34	0.37	
0.35	0.38	
0.35	0.39	
0.36	0.40	

Number of values= 25

Mean of values 0.37
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

31-33	****
34-36	*****
37-39	*****
40-42	****
43-45	

VITRINITE REFLECTANCE VALUES

Well Name: DUTSON DOWNS-1
Depth: 5360 ft

Sorted List

0.28	0.34
0.30	0.35
0.31	0.36
0.31	0.36
0.31	0.37
0.31	0.37
0.33	0.37
0.33	0.42
0.33	0.45
0.33	0.47

Number of values= 20

Mean of values 0.35
Standard Deviation 0.05

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

28-30	**
31-33	*****
34-36	***
37-39	**
40-42	*
43-45	*
46-48	*

VITRINITE REFLECTANCE VALUES

Well Name: DUTSON DOWNS-1
Depth: 6110 ft

Sorted List

0.34	0.44	0.51
0.36	0.45	0.52
0.37	0.45	0.52
0.38	0.45	0.53
0.38	0.46	
0.39	0.47	
0.39	0.48	
0.40	0.48	
0.41	0.49	
0.44	0.51	

Number of values= 24

Mean of values 0.44
Standard Deviation 0.06

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

34-36	**
37-39	*****
40-42	**
43-45	*****
46-48	****
49-51	***
52-54	***

VITRINITE REFLECTANCE VALUES

Well Name: MERRIMAN-1
Depth: 2210 ft

Sorted List

0.21
0.22
0.32
0.33
0.34

Number of values= 5

Mean of values 0.28
Standard Deviation 0.06

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

21-23 **
24-26
27-29
30-32 *
33-35 **

VITRINITE REFLECTANCE VALUES

Well Name: MERRIMAN-1
Depth: 2510 ft

Sorted List

0.20	0.24	0.27	0.31
0.20	0.25	0.28	
0.21	0.25	0.28	
0.23	0.25	0.28	
0.23	0.25	0.29	
0.23	0.26	0.29	
0.23	0.26	0.29	
0.24	0.26	0.29	
0.24	0.26	0.31	
0.24	0.26	0.31	

Number of values= 31

Mean of values 0.26
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

20-22	***
23-25	*****
26-28	*****
29-31	*****

VITRINITE REFLECTANCE VALUES

Well Name: MERRIMAN-1
Depth: 4350 ft

Sorted List

0.24	0.30	0.36
0.26	0.30	0.36
0.26	0.31	0.36
0.27	0.32	0.37
0.28	0.32	0.39
0.28	0.32	0.40
0.29	0.32	
0.29	0.33	
0.29	0.34	
0.30	0.36	

Number of values= 26

Mean of values 0.32
Standard Deviation 0.04

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

24-26	***
27-29	*****
30-32	*****
33-35	**
36-38	****
39-41	

VITRINITE REFLECTANCE VALUES

Well Name: MERRIMAN-1
Depth: 4720 ft

Sorted List

0.33	0.38	0.41
0.36	0.39	0.41
0.36	0.39	0.43
0.36	0.39	0.45
0.37	0.39	
0.37	0.39	
0.37	0.39	
0.38	0.40	
0.38	0.40	
0.38	0.40	

Number of values= 24

Mean of values 0.39
Standard Deviation 0.02

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

33-35	*
36-38	*****
39-41	*****
42-44	*
45-47	*

VITRINITE REFLECTANCE VALUES

Well Name: MERRIMAN-1
Depth: 5081 ft

Sorted List

0.41
0.47

Number of values= 2

Mean of values 0.44
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

41-43 *
44-46
47-49 *

VITRINITE REFLECTANCE VALUES

Well Name: MERRIMAN-1
Depth: 5960 ft

Sorted List

0.40 0.53
0.40 0.55
0.41
0.42
0.44
0.44
0.46
0.49
0.50
0.51

Number of values= 12

Mean of values 0.46
Standard Deviation 0.05

HISTOGRAM OF VALUES Reflectance values multiplied by 100

40-42	****
43-45	**
46-48	*
49-51	***
52-54	*
55-57	*

VITRINITE REFLECTANCE VALUES

Well Name: NORTH SEASPRAY-1
Depth: 1910 FT

Sorted List

0.25	0.28	0.31
0.26	0.29	0.31
0.27	0.29	
0.27	0.29	
0.27	0.29	
0.27	0.29	
0.28	0.29	
0.28	0.29	
0.28	0.30	
0.28	0.30	

Number of values= 22

Mean of values 0.28
Standard Deviation 0.01

HISTOGRAM OF VALUES Reflectance values multiplied by 100

25-27	**
28-30	*****
31-33	**

VITRINITE REFLECTANCE VALUES

Well Name: NORTH SEASPRAY-1
Depth: 2100 ft

Sorted List

0.26	0.28	0.31
0.26	0.29	0.32
0.26	0.29	0.33
0.26	0.29	0.33
0.26	0.30	
0.26	0.30	
0.28	0.30	
0.28	0.31	
0.28	0.31	
0.28	0.31	

Number of values= 24

Mean of values 0.29
Standard Deviation 0.02

HISTOGRAM OF VALUES
Reflectance values multiplied by 100

26-28	*****
29-31	*****
32-34	***

VITRINITE REFLECTANCE VALUES

Well Name: NORTH SEASPRAY-1
Depth: 3310 ft

Sorted List

0.28	0.33	0.37
0.30	0.34	0.38
0.30	0.34	0.44
0.30	0.34	0.44
0.30	0.34	
0.31	0.34	
0.31	0.35	
0.31	0.35	
0.32	0.36	
0.32	0.36	

Number of values= 24

Mean of values 0.34
Standard Deviation 0.04

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

28-30	*****
31-33	*****
34-36	*****
37-39	**
40-42	
43-45	**

VITRINITE REFLECTANCE VALUES

Well Name: NORTH SEASPRAY-1
Depth: 3504 ft

Sorted List

0.28	0.32	0.35
0.28	0.32	0.36
0.29	0.32	0.37
0.30	0.33	0.37
0.30	0.33	
0.30	0.34	
0.30	0.34	
0.31	0.34	
0.31	0.35	
0.31	0.35	

Number of values= 24

Mean of values 0.32
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

28-30	*****
31-33	*****
34-36	*****
37-39	**

VITRINITE REFLECTANCE VALUES

Well Name: NORTH SEASPRAY-1
Depth: 3690 ft

Sorted List

0.25	0.31	0.37
0.28	0.31	0.40
0.28	0.31	0.40
0.28	0.32	0.40
0.29	0.32	
0.29	0.32	
0.30	0.33	
0.30	0.33	
0.30	0.34	
0.30	0.36	

Number of values= 24

Mean of values 0.32
Standard Deviation 0.04

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

25-27	*
28-30	*****
31-33	*****
34-36	**
37-39	*
40-42	***

VITRINITE REFLECTANCE VALUES

Well Name: NORTH SEASPRAY-1
Depth: 3800 ft

Sorted List

0.25	0.32	0.36
0.27	0.33	0.36
0.29	0.33	0.40
0.30	0.33	0.40
0.30	0.34	
0.31	0.34	
0.31	0.34	
0.31	0.35	
0.32	0.35	
0.32	0.36	

Number of values= 24

Mean of values 0.33
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

25-27	*
28-30	****
31-33	*****
34-36	*****
37-39	
40-42	**

VITRINITE REFLECTANCE VALUES

Well Name: NORTH SEASPRAY-1
Depth: 4000 ft

Sorted List

0.31
0.33
0.34
0.35
0.36
0.41
0.42
0.43
0.45
0.45

Number of values= 10

Mean of values 0.39
Standard Deviation 0.05

HISTOGRAM OF VALUES
Reflectance values multiplied by 100

31-33 **
34-36 ***
37-39
40-42 **
43-45 ***

VITRINITE REFLECTANCE VALUES

Well Name: NORTH SEASPRAY-1
Depth: 5000 ft

Sorted List

0.29 0.42
0.35 0.42
0.35 0.42
0.35 0.43
0.35 0.45
0.36 0.48
0.37
0.38
0.38
0.40

Number of values= 16

Mean of values 0.39
Standard Deviation 0.05

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

29-31	*
32-34	
35-37	*****
38-40	***
41-43	****
44-46	*
47-49	*

VITRINITE REFLECTANCE VALUES

Well Name: SEASPRAY-1
Depth: 4466 ft

Sorted List

0.29	0.36	0.42
0.30	0.37	0.44
0.32	0.37	
0.32	0.38	
0.32	0.38	
0.33	0.39	
0.33	0.39	
0.33	0.40	
0.35	0.41	
0.36	0.41	

Number of values= 22

Mean of values 0.36
Standard Deviation 0.04

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

29-31	**
32-34	*****
35-37	****
38-40	****
41-43	***
44-46	*

VITRINITE REFLECTANCE VALUES

Well Name: SEASPRAY-1
Depth: 4730 ft

Sorted List

0.28	0.34	0.44
0.29	0.34	0.48
0.29	0.37	0.49
0.30	0.37	
0.31	0.39	
0.32	0.39	
0.33	0.39	
0.33	0.43	
0.34	0.43	
0.34	0.43	

Number of values= 23

Mean of values 0.37
Standard Deviation 0.06

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

28-30	****
31-33	****
34-36	****
37-39	*****
40-42	
43-45	****
46-48	*
49-51	*

VITRINITE REFLECTANCE VALUES

Well Name: SEASPRAY-1
Depth: 5000 ft

Sorted List

0.33 0.38
0.34 0.38
0.35 0.38
0.35 0.39
0.36 0.40
0.36 0.41
0.37
0.37
0.37
0.38

Number of values= 16

Mean of values 0.37
Standard Deviation 0.02

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

33-35 ****
36-38 *****
39-41 ***

VITRINITE REFLECTANCE VALUES

Well Name: WELLINGTON PARK-1
Depth: 1880 FT

Sorted List

0.27
0.29
0.31
0.34
0.35
0.38
0.39
0.40
0.50

Number of values= 9

Mean of values 0.36
Standard Deviation 0.07

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

27-29 **
30-32 *
33-35 **
36-38 *
39-41 **
42-44
45-47
48-50 *

VITRINITE REFLECTANCE VALUES

Well Name: WELLINGTON PARK-1
Depth: 3310 ft

Sorted List

0.24	0.27	0.31
0.25	0.27	0.32
0.25	0.28	0.33
0.25	0.29	0.33
0.26	0.29	
0.26	0.29	
0.26	0.31	
0.27	0.31	
0.27	0.31	
0.27	0.31	

Number of values= 24

Mean of values 0.28
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

24-26	*****
27-29	*****
30-32	*****
33-35	**

VITRINITE REFLECTANCE VALUES

Well Name: WELLINGTON PARK-1
Depth: 3650 ft

Sorted List

0.28 0.35
0.28 0.35
0.32 0.35
0.32 0.35
0.33 0.36
0.33 0.40
0.33
0.34
0.34
0.34

Number of values= 16

Mean of values 0.34
Standard Deviation 0.03

HISTOGRAM OF VALUES
Reflectance values multiplied by 100

28-30 **
31-33 *****
34-36 *****
37-39
40-42 *

VITRINITE REFLECTANCE VALUES

Well Name: WELLINGTON PARK-1
Depth: 3830 ft

Sorted List

0.28	0.33	0.39
0.28	0.34	0.39
0.29	0.34	
0.29	0.35	
0.30	0.35	
0.30	0.36	
0.30	0.36	
0.31	0.36	
0.31	0.36	
0.32	0.38	

Number of values= 22

Mean of values 0.33
Standard Deviation 0.03

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

28-30	*****
31-33	***
34-36	*****
37-39	***

VITRINITE REFLECTANCE VALUES

Well Name: WELLINGTON PARK-1
Depth: 7376 ft

Sorted List

0.46	0.62	0.66
0.49	0.63	0.66
0.55	0.63	0.66
0.56	0.63	0.67
0.57	0.63	0.67
0.59	0.63	0.68
0.59	0.64	0.72
0.62	0.65	0.72
0.62	0.65	0.76
0.62	0.66	0.77

Number of values= 30

Mean of values 0.63
Standard Deviation 0.07

HISTOGRAM OF VALUES

Reflectance values multiplied by 100

46-48	*
49-51	*
52-54	
55-57	***
58-60	**
61-63	*****
64-66	*****
67-69	*
70-72	
73-75	
76-78	**

Enclosure 1:

Cross Section - Lake Wellington Park
to Merriman-1

PE 801905

801903 082
C84

DEPT. PRIMARY INDUSTRIES
PETROLEUM DEVELOPMENT
THIS DOCUMENT HAS BEEN
SCANNED
ON OR BEFORE JANUARY 2004

PE801905

This is an enclosure indicator page.
The enclosure PE801905 is enclosed within the
container PE801903 at this location in this
document.

The enclosure PE801905 has the following characteristics:

ITEM_BARCODE = PE801905
CONTAINER_BARCODE = PE801903
NAME = PEP 109 Stratigraphic Cross-Section
BASIN = GIPPSLAND
ONSHORE? = Y
DATA_TYPE = WELL
DATA_SUB_TYPE = WELL_CORRELATION
DESCRIPTION = PEP 109 Stratigraphic Cross-Section,
Merriman-1 to Wellington Park-1
(enclosure 1 from Palynology and Source
Rock Study of PEP 109), Onshore
Gippsland Basin, Cluff Resources
Pacific Ltd, April 1991
REMARKS =
DATE_WRITTEN = 30-APR-1991
DATE_PROCESSED =
DATE_RECEIVED =
RECEIVED_FROM = Cluff Resources Pacific Ltd
WELL_NAME = Wellington Park-1
CONTRACTOR =
AUTHOR =
ORIGINATOR =
TOP_DEPTH =
BOTTOM_DEPTH =
ROW_CREATED_BY = FH11_SW

(Inserted by DNRE - Vic Govt Mines Dept)

Enclosure 2:

Cross Section - Golden Beach 1A to
Colliers Hill

PE 801908.

801903 ~~085~~ 085

DEPT. PRIMARY INDUSTRIES
PETROLEUM DEVELOPMENT
THIS DOCUMENT HAS BEEN
SCANNED
ON OR BEFORE JANUARY 2004

PE801908

This is an enclosure indicator page.
The enclosure PE801908 is enclosed within the
container PE801903 at this location in this
document.

The enclosure PE801908 has the following characteristics:

ITEM_BARCODE = PE801908
CONTAINER_BARCODE = PE801903
NAME = PEP 109 Stratigraphic Cross-Section
BASIN = GIPPSLAND
ONSHORE? = Y
DATA_TYPE = WELL
DATA_SUB_TYPE = WELL_CORRELATION
DESCRIPTION = PEP 109 Stratigraphic Cross-Section,
Colliers Hill-1 to Golden Beach-1A
(enclosure 2 from Palynology and Source
Rock Study of PEP 109), Onshore
Gippsland Basin, Cluff Resources
Pacific Ltd, April 1991
REMARKS =
DATE_WRITTEN = 30-APR-1991
DATE_PROCESSED =
DATE_RECEIVED =
RECEIVED_FROM = Cluff Resources Pacific Ltd
WELL_NAME = Golden Beach-1A
CONTRACTOR =
AUTHOR =
ORIGINATOR =
TOP_DEPTH =
BOTTOM_DEPTH =
ROW_CREATED_BY = FH11_SW

(Inserted by DNRE - Vic Govt Mines Dept)